

# Chapter 1 Relations and Functions

## EXERCISE 1.1

### Question 1:

Determine whether each of the following relations are reflexive, symmetric and transitive.

- (i) Relation  $R$  in the set  $A = \{1, 2, 3, \dots, 13, 14\}$  defined as

$$R = \{(x, y) : 3x - y = 0\}$$

- (ii) Relation  $R$  in the set of  $\mathbb{N}$  natural numbers defined as

$$R = \{(x, y) : y = x + 5 \text{ and } x < 4\}$$

- (iii) Relation  $R$  in the set  $A = \{1, 2, 3, 4, 5, 6\}$  defined as

$$R = \{(x, y) : y \text{ is divisible by } x\}$$

- (iv) Relation  $R$  in the set of  $\mathbb{Z}$  integers defined as

$$R = \{(x, y) : x - y \text{ is an integer}\}$$

- (v) Relation  $R$  in the set of human beings in a town at a particular time given by

(a)  $R = \{(x, y) : x \text{ and } y \text{ work at the same place}\}$

(b)  $R = \{(x, y) : x \text{ and } y \text{ live in the same locality}\}$

(c)  $R = \{(x, y) : x \text{ is exactly 7cm taller than } y\}$

(d)  $R = \{(x, y) : x \text{ is wife of } y\}$

(e)  $R = \{(x, y) : x \text{ is father of } y\}$

### Solution:

(i)  $R = \{(1, 3), (2, 6), (3, 9), (4, 12)\}$

$R$  is not reflexive because  $(1, 1), (2, 2), \dots$  and  $(14, 14) \notin R$ .

$R$  is not symmetric because  $(1, 3) \in R$ , but  $(3, 1) \notin R$ . [since  $3(3) \neq 0$ ].

$R$  is not transitive because  $(1, 3), (3, 9) \in R$ , but  $(1, 9) \notin R$ . [ $3(1) - 9 \neq 0$ ].

Hence,  $R$  is neither reflexive nor symmetric nor transitive.

(ii)  $R = \{(1, 6), (2, 7), (3, 8)\}$

$R$  is not reflexive because  $(1, 1) \notin R$ .

$R$  is not symmetric because  $(1, 6) \in R$  but  $(6, 1) \notin R$ .

$R$  is not transitive because there isn't any ordered pair in  $R$  such that  $(x, y), (y, z) \in R$ , so  $(x, z) \notin R$ .

Hence,  $R$  is neither reflexive nor symmetric nor transitive.

(iii)  $R = \{(x, y) : y \text{ is divisible by } x\}$

We know that any number other than 0 is divisible by itself.

Thus,  $(x, x) \in R$

So,  $R$  is reflexive.

$(2,4) \in R$  [because 4 is divisible by 2]

But  $(4,2) \notin R$  [since 2 is not divisible by 4]

So,  $R$  is not symmetric.

Let  $(x,y)$  and  $(y,z) \in R$ . So,  $y$  is divisible by  $x$  and  $z$  is divisible by  $y$ .

So,  $z$  is divisible by  $x \Rightarrow (x,z) \in R$

So,  $R$  is transitive.

So,  $R$  is reflexive and transitive but not symmetric.

(iv)  $R = \{(x,y) : x - y \text{ is an integer}\}$

For  $x \in \mathbb{Z}$ ,  $(x,x) \in R$  because  $x - x = 0$  is an integer.

So,  $R$  is reflexive.

For,  $x, y \in \mathbb{Z}$ , if  $(x,y) \in R$ , then  $x - y$  is an integer  $\Rightarrow (y - x)$  is an integer.

So,  $(y,x) \in R$

So,  $R$  is symmetric.

Let  $(x,y)$  and  $(y,z) \in R$ , where  $x, y, z \in \mathbb{Z}$ .

$\Rightarrow (x - y)$  and  $(y - z)$  are integers.

$\Rightarrow x - z = (x - y) + (y - z)$  is an integer.

So,  $R$  is transitive.

So,  $R$  is reflexive, symmetric and transitive.

(v)

a)  $R = \{(x,y) : x \text{ and } y \text{ work at the same place}\}$

$R$  is reflexive because  $(x,x) \in R$

$R$  is symmetric because,

If  $(x,y) \in R$ , then  $x$  and  $y$  work at the same place and  $y$  and  $x$  also work at the same place.  $(y,x) \in R$ .

$R$  is transitive because,

Let  $(x,y), (y,z) \in R$

$x$  and  $y$  work at the same place and  $y$  and  $z$  work at the same place.

Then,  $x$  and  $z$  also works at the same place.  $(x,z) \in R$ .

Hence,  $R$  is reflexive, symmetric and transitive.

b)  $R = \{(x,y) : x \text{ and } y \text{ live in the same locality}\}$

$R$  is reflexive because  $(x,x) \in R$

$R$  is symmetric because,

If  $(x,y) \in R$ , then  $x$  and  $y$  live in the same locality and  $y$  and  $x$  also live in the same locality  $(y,x) \in R$ .

$R$  is transitive because,

Let  $(x, y), (y, z) \in R$

$x$  and  $y$  live in the same locality and  $y$  and  $z$  live in the same locality.

Then  $x$  and  $z$  also live in the same locality.  $(x, z) \in R$ .

Hence,  $R$  is reflexive, symmetric and transitive.

c)  $R = \{(x, y) : x \text{ is exactly } 7\text{cm taller than } y\}$

$R$  is not reflexive because  $(x, x) \notin R$ .

$R$  is not symmetric because,

If  $(x, y) \in R$ , then  $x$  is exactly  $7\text{cm}$  taller than  $y$  and  $y$  is clearly not taller than  $x$ .

$(y, x) \notin R$ .

$R$  is not transitive because,

Let  $(x, y), (y, z) \in R$

$x$  is exactly  $7\text{cm}$  taller than  $y$  and  $y$  is exactly  $7\text{cm}$  taller than  $z$ .

Then  $x$  is exactly  $14\text{cm}$  taller than  $z$ .  $(x, z) \notin R$ .

Hence,  $R$  is neither reflexive nor symmetric nor transitive.

d)  $R = \{(x, y) : x \text{ is wife of } y\}$

$R$  is not reflexive because  $(x, x) \notin R$ .

$R$  is not symmetric because,

Let  $(x, y) \in R$ ,  $x$  is the wife of  $y$  and  $y$  is not the wife of  $x$ .  $(y, x) \notin R$ .

$R$  is not transitive because,

Let  $(x, y), (y, z) \in R$

$x$  is wife of  $y$  and  $y$  is wife of  $z$ , which is not possible.

$(x, z) \notin R$ .

Hence,  $R$  is neither reflexive nor symmetric nor transitive.

e)  $R = \{(x, y) : x \text{ is father of } y\}$

$R$  is not reflexive because  $(x, x) \notin R$ .

$R$  is not symmetric because,

Let  $(x, y) \in R$ ,  $x$  is the father of  $y$  and  $y$  is not the father of  $x$ .  $(y, x) \notin R$ .

$R$  is not transitive because,

Let  $(x, y), (y, z) \in R$

$x$  is father of  $y$  and  $y$  is father of  $z$ ,  $x$  is not father of  $z$ .  $(x, z) \notin R$ .

Hence,  $R$  is neither reflexive nor symmetric nor transitive.

### Question 2:

Show that the relation  $R$  in the set  $\mathbb{R}$  of real numbers, defined as  $R = \{(a,b) : a \leq b^2\}$  is neither reflexive nor symmetric nor transitive.

#### Solution:

$$R = \{(a,b) : a \leq b^2\}$$

$$\left(\frac{1}{2}, \frac{1}{2}\right) \notin R \quad \text{because} \quad \frac{1}{2} > \left(\frac{1}{2}\right)^2$$

$R$  is not reflexive.

$(1,4) \in R$  as  $1 < 4$ . But  $4$  is not less than  $1^2$ .

$$(4,1) \notin R$$

$R$  is not symmetric.

$$(3,2)(2,1.5) \in R \quad [\text{Because } 3 < 2^2=4 \text{ and } 2 < (1.5)^2=2.25]$$

$$3 > (1.5)^2 = 2.25$$

$$\therefore (3,1.5) \notin R$$

$R$  is not transitive.

$R$  is neither reflexive nor symmetric nor transitive.

### Question 3:

Check whether the relation  $R$  defined in the set  $\{1,2,3,4,5,6\}$  as  $R = \{(a,b) : b = a + 1\}$  is reflexive, symmetric or transitive.

#### Solution:

$$A = \{1,2,3,4,5,6\}$$

$$R = \{(a,b) : b = a + 1\}$$

$$R = \{(1,2), (2,3), (3,4), (4,5), (5,6)\}$$

$$(a,a) \notin R, a \in A$$

$$(1,1), (2,2), (3,3), (4,4), (5,5) \notin R$$

$R$  is not reflexive.

$$(1,2) \in R, \text{ but } (2,1) \notin R$$

R is not symmetric.

$$(1,2), (2,3) \in R$$

$$(1,3) \notin R$$

R is not transitive.

R is neither reflexive nor symmetric nor transitive.

#### Question 4:

Show that the relation R in R defined as  $R = \{(a,b) : a \leq b\}$  is reflexive and transitive, but not symmetric.

#### Solution:

$$R = \{(a,b) : a \leq b\}$$

$$(a,a) \in R$$

R is reflexive.

$$(2,4) \in R \text{ (as } 2 < 4)$$

$$(4,2) \notin R \text{ (as } 4 > 2)$$

R is not symmetric.

$$(a,b), (b,c) \in R$$

$$a \leq b \text{ and } b \leq c$$

$$\Rightarrow a \leq c$$

$$\Rightarrow (a,c) \in R$$

R is transitive.

R is reflexive and transitive but not symmetric.

#### Question 5:

Check whether the relation R in R defined as  $R = \{(a,b) : a \leq b^3\}$  is reflexive, symmetric or transitive.

#### Solution:

$$R = \{(a,b) : a \leq b^3\}$$

$$\left(\frac{1}{2}, \frac{1}{2}\right) \notin R, \text{ since } \frac{1}{2} > \left(\frac{1}{2}\right)^3$$

R is not reflexive.

$$(1,2) \in R \text{ (as } 1 < 2^3 = 8)$$

$$(2,1) \notin R \text{ (as } 2^3 > 1 = 8)$$

R is not symmetric.

$$\left(3, \frac{3}{2}\right), \left(\frac{3}{2}, \frac{6}{5}\right) \in R, \text{ since } 3 < \left(\frac{3}{2}\right)^3 \text{ and } \frac{2}{3} < \left(\frac{6}{5}\right)^3$$

$$\left(3, \frac{6}{5}\right) \notin R \text{ as } 3 > \left(\frac{6}{5}\right)^3$$

R is not transitive.

R is neither reflexive nor symmetric nor transitive.

### Question 6:

Show that the relation R in the set  $\{1,2,3\}$  given by  $R = \{(1,2), (2,1)\}$  is symmetric but neither reflexive nor transitive.

### Solution:

$$A = \{1,2,3\}$$

$$R = \{(1,2), (2,1)\}$$

$$(1,1), (2,2), (3,3) \notin R$$

R is not reflexive.

$$(1,2) \in R \text{ and } (2,1) \in R$$

R is symmetric.

$$(1,2) \in R \text{ and } (2,1) \in R$$

$$(1,1) \notin R$$

R is not transitive.

R is symmetric, but not reflexive or transitive.

### Question 7:

Show that the relation R in the set A of all books in a library of a college, given by  $R = \{(x,y) : x \text{ and } y \text{ have same number of pages}\}$  is an equivalence relation.

### Solution:

$$R = \{(x,y) : x \text{ and } y \text{ have same number of pages}\}$$

R is reflexive since  $(x,x) \in R$  as  $x$  and  $x$  have same number of pages.

R is reflexive.

$$(x, y) \in R$$

$x$  and  $y$  have same number of pages and  $y$  and  $x$  have same number of pages  $(y, x) \in R$

R is symmetric.

$$(x, y) \in R, (y, z) \in R$$

$x$  and  $y$  have same number of pages,  $y$  and  $z$  have same number of pages.

Then  $x$  and  $z$  have same number of pages.

$$(x, z) \in R$$

R is transitive.

R is an equivalence relation.

### Question 8:

Show that the relation R in the set  $A = \{1, 2, 3, 4, 5\}$  given by  $R = \{(a, b) : |a - b| \text{ is even}\}$  is an equivalence relation. Show that all the elements of  $\{1, 3, 5\}$  are related to each other and all the elements of  $\{2, 4\}$  are related to each other. But no element of  $\{1, 3, 5\}$  is related to any element of  $\{2, 4\}$ .

### Solution:

$$a \in A$$

$$|a - a| = 0 \text{ (which is even)}$$

R is reflexive.

$$(a, b) \in R$$

$$\Rightarrow |a - b| \text{ [is even]}$$

$$\Rightarrow |-(a - b)| = |b - a| \text{ [is even]}$$

$$(b, a) \in R$$

R is symmetric.

$$(a, b) \in R \text{ and } (b, c) \in R$$

$$\Rightarrow |a - b| \text{ is even and } |b - c| \text{ is even}$$

$$\Rightarrow (a - b) \text{ is even and } (b - c) \text{ is even}$$

$$\Rightarrow (a - c) = (a - b) + (b - c) \text{ is even}$$

$\Rightarrow |a - b|$  is even

$\Rightarrow (a, c) \in R$

R is transitive.

R is an equivalence relation.

All elements of  $\{1, 3, 5\}$  are related to each other because they are all odd. So, the modulus of the difference between any two elements is even.

Similarly, all elements  $\{2, 4\}$  are related to each other because they are all even.

No element of  $\{1, 3, 5\}$  is related to any elements of  $\{2, 4\}$  as all elements of  $\{1, 3, 5\}$  are odd and all elements of  $\{2, 4\}$  are even. So, the modulus of the difference between the two elements will not be even.

### Question 9:

Show that each of the relation R in the set  $A = \{x \in Z : 0 \leq x \leq 12\}$ , given by

i.  $R = \{(a, b) : |a - b| \text{ is a multiple of } 4\}$

ii.  $R = \{(a, b) : a = b\}$

Is an equivalence relation. Find the set of all elements related to 1 in each case.

### Solution:

$$A = \{x \in Z : 0 \leq x \leq 12\} = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$$

i.  $R = \{(a, b) : |a - b| \text{ is a multiple of } 4\}$

$$a \in A, (a, a) \in R \quad [ |a - a| = 0 \text{ is a multiple of } 4 ]$$

R is reflexive.

$$(a, b) \in R \Rightarrow |a - b| \text{ [is a multiple of } 4]$$

$$\Rightarrow |-(a - b)| = |b - a| \text{ [is a multiple of } 4]$$

$$(b, a) \in R$$

R is symmetric.

$$(a, b) \in R \text{ and } (b, c) \in R$$

$$\Rightarrow |a - b| \text{ is a multiple of } 4 \text{ and } |b - c| \text{ is a multiple of } 4$$

$$\Rightarrow (a - b) \text{ is a multiple of } 4 \text{ and } (b - c) \text{ is a multiple of } 4$$

$$\Rightarrow (a - c) = (a - b) + (b - c) \text{ is a multiple of } 4$$

$$\Rightarrow |a - c| \text{ is a multiple of } 4$$

$$\Rightarrow (a, c) \in R$$

R is transitive.

R is an equivalence relation.

The set of elements related to 1 is  $\{1, 5, 9\}$  as

$$|1-1| = 0 \text{ is a multiple of 4.}$$

$$|5-1| = 4 \text{ is a multiple of 4.}$$

$$|9-1| = 8 \text{ is a multiple of 4.}$$

ii.  $R = \{(a, b) : a = b\}$

$$a \in A, (a, a) \in R \quad [\text{since } a=a]$$

R is reflexive.

$$(a, b) \in R$$

$$\Rightarrow a = b$$

$$\Rightarrow b = a$$

$$\Rightarrow (b, a) \in R$$

R is symmetric.

$$(a, b) \in R \text{ and } (b, c) \in R$$

$$\Rightarrow a = b \text{ and } b = c$$

$$\Rightarrow a = c$$

$$\Rightarrow (a, c) \in R$$

R is transitive.

R is an equivalence relation.

The set of elements related to 1 is  $\{1\}$ .

### Question 10:

Give an example of a relation, which is

- i. Symmetric but neither reflexive nor transitive.
- ii. Transitive but neither reflexive nor symmetric.
- iii. Reflexive and symmetric but not transitive.
- iv. Reflexive and transitive but not symmetric.
- v. Symmetric and transitive but not reflexive.

### Solution:

- i.

$$A = \{5, 6, 7\}$$

$$R = \{(5, 6), (6, 5)\}$$

$$(5, 5), (6, 6), (7, 7) \notin R$$

R is not reflexive as  $(5, 5), (6, 6), (7, 7) \notin R$

$(5, 6), (6, 5) \in R$  and  $(6, 5) \in R$ , R is symmetric.

$\Rightarrow (5, 6), (6, 5) \in R$ , but  $(5, 5) \notin R$

R is not transitive.

Relation R is symmetric but not reflexive or transitive.

ii.  $R = \{(a, b) : a < b\}$

$a \in R, (a, a) \notin R$  [since  $a$  cannot be less than itself]

R is not reflexive.

$$(1, 2) \in R \text{ (as } 1 < 2)$$

But 2 is not less than 1

$$\therefore (2, 1) \notin R$$

R is not symmetric.

$$(a, b), (b, c) \in R$$

$$\Rightarrow a < b \text{ and } b < c$$

$$\Rightarrow a < c$$

$$\Rightarrow (a, c) \in R$$

R is transitive.

Relation R is transitive but not reflexive and symmetric.

iii.  $A = \{4, 6, 8\}$

$$A = \{(4, 4), (6, 6), (8, 8), (4, 6), (6, 8), (8, 6)\}$$

R is reflexive since  $a \in A, (a, a) \in R$

R is symmetric since  $(a, b) \in R$

$$\Rightarrow (b, a) \in R \text{ for } a, b \in R$$

R is not transitive since  $(4, 6), (6, 8) \in R$ , but  $(4, 8) \notin R$

R is reflexive and symmetric but not transitive.

iv.  $R = \{(a, b) : a^3 > b^3\}$

$$(a, a) \in R$$

R is reflexive.

$$(2, 1) \in R$$

$$\text{But } (1, 2) \notin R$$

$\therefore R$  is not symmetric.

$$(a, b), (b, c) \in R$$

$$\Rightarrow a^3 \geq b^3 \text{ and } b^3 < c^3$$

$$\Rightarrow a^3 < c^3$$

$$\Rightarrow (a, c) \in R$$

$\therefore R$  is transitive.

$R$  is reflexive and transitive but not symmetric

v.

$$A = \{1, 3, 5\}$$

Define a Relation  $R$

On  $A$ .

$$R : A \rightarrow A$$

$$R = \{(1,3) (3,1) (1,1) (3,3)\}$$

Relation  $R$  is not Reflexive as  $(5, 5) \notin R$

Relation  $R$  is Symmetric as

$$(1,3) \in R \Rightarrow (3,1) \in R$$

Relation  $R$  is Transitive

$$(a, b) \in R, (b, c) \in R \Rightarrow (a, c) \in R$$

$$(3,1) \in R, (1,1) \in R \Rightarrow (3,1) \in R$$

**Alternative Answer**

$R = (a, b) : a$  is brother of  $b$  {suppose  $a$  and  $b$  are male}

$Ref \rightarrow a$  is not brother of  $a$

So,  $(a, a) \notin R$

Relation  $R$  is not Reflexive

Symmetric  $\rightarrow a$  is brother of  $b$  so

$b$  is brother of  $a$

$$a, b \in R \Rightarrow (b, a) \in R$$

Transitive  $\rightarrow a$  is brother of  $b$  and

$b$  is brother of  $c$  so

$a$  is brother of  $c$

$$(a, b) \in R, (b, c) \in R \Rightarrow (a, c) \in R$$

### Question 11:

Show that the relation  $R$  in the set  $A$  of points in a plane given by

$$R = \{(P, Q) : \text{Distance of the point } P \text{ from the origin is same as the distance of the point } Q \text{ from the origin}\}$$

, is an equivalence relation. Further, show that the set of all points related to a point  $P \neq (0,0)$  is the circle passing through  $P$  with origin as centre.

### Solution:

$$R = \{(P, Q) : \text{Distance of the point } P \text{ from the origin is same as the distance of the point } Q \text{ from the origin}\}$$

Clearly,  $(P, P) \in R$

$R$  is reflexive.

$$(P, Q) \in R$$

Clearly  $R$  is symmetric.

$$(P, Q), (Q, S) \in R$$

$\Rightarrow$  The distance of  $P$  and  $Q$  from the origin is the same and also, the distance of  $Q$  and  $S$  from the origin is the same.

$\Rightarrow$  The distance of  $P$  and  $S$  from the origin is the same.

$$(P, S) \in R$$

$R$  is transitive.

$R$  is an equivalence relation.

The set of points related to  $P \neq (0,0)$  will be those points whose distance from origin is same as distance of  $P$  from the origin.

Set of points forms a circle with the centre as origin and this circle passes through  $P$ .

### Question 12:

Show that the relation  $R$  in the set  $A$  of all triangles as  $R = \{(T_1, T_2) : T_1 \text{ is similar to } T_2\}$ , is an equivalence relation. Consider three right angle triangles  $T_1$  with sides 3,4,5,  $T_2$  with sides 5,12,13 and  $T_3$  with sides 6,8,10. Which triangle among  $T_1, T_2, T_3$  are related?

### Solution:

$$R = \{(T_1, T_2) : T_1 \text{ is similar to } T_2\}$$

$R$  is reflexive since every triangle is similar to itself.

If  $(T_1, T_2) \in R$ , then  $T_1$  is similar to  $T_2$ .

$T_2$  is similar to  $T_1$ .

$$\Rightarrow (T_2, T_1) \in R$$

$R$  is symmetric.

$$(T_1, T_2), (T_2, T_3) \in R$$

is similar to  $T_2$  and  $T_2$  is similar to  $T_3$ .

$\therefore T_1$  is similar to  $T_3$ .

$\Rightarrow (T_1, T_3) \in R$

$\therefore R$  is transitive.

$$\frac{3}{6} = \frac{4}{8} = \frac{5}{10} = \left(\frac{1}{2}\right)$$

$\therefore$  Corresponding sides of triangles  $T_1$  and  $T_3$  are in the same ratio.

Triangle  $T_1$  is similar to triangle  $T_3$ .

Hence,  $T_1$  is related to  $T_3$ .

### Question 13:

Show that the relation  $R$  in the set  $A$  of all polygons as  $R = \{(P_1, P_2) : P_1 \text{ and } P_2 \text{ have same number of sides}\}$ , is an equivalence relation. What is the set of all elements in  $A$  related to the right angle triangle  $T$  with sides 3, 4 and 5?

**Solution:**

$$R = \{(P_1, P_2) : P_1 \text{ and } P_2 \text{ have same number of sides}\}$$

$(P_1, P_2) \in R$  as same polygon has same number of sides.

$\therefore R$  is reflexive.

$$(P_1, P_2) \in R$$

$\Rightarrow P_1$  and  $P_2$  have same number of sides.

$\Rightarrow P_2$  and  $P_1$  have same number of sides.

$$\Rightarrow (P_2, P_1) \in R$$

$\therefore R$  is symmetric.

$$(P_1, P_2), (P_2, P_3) \in R$$

$\Rightarrow P_1$  and  $P_2$  have same number of sides.

$P_2$  and  $P_3$  have same number of sides.

$\Rightarrow P_1$  and  $P_3$  have same number of sides.

$$\Rightarrow (P_1, P_3) \in R$$

$\therefore R$  is transitive.

$R$  is an equivalence relation.

The elements in  $A$  related to right-angled triangle (T) with sides 3, 4, 5 are those polygons which have three sides.

Set of all elements in  $A$  related to triangle T is the set of all triangles.

**Question 14:**

Let  $L$  be the set of all lines in XY plane and  $R$  be the relation in  $L$  defined as

$R = \{(L_1, L_2) : L_1 \text{ is parallel to } L_2\}$ . Show that  $R$  is an equivalence relation. Find the set of all lines related to the line  $y = 2x + 4$ .

**Solution:**

$$R = \{(L_1, L_2) : L_1 \text{ is parallel to } L_2\}$$

$R$  is reflexive as any line  $L_1$  is parallel to itself i.e.,  $(L_1, L_1) \in R$

If  $(L_1, L_2) \in R$ , then

$\Rightarrow L_1$  is parallel to  $L_2$ .

$\Rightarrow L_2$  is parallel to  $L_1$ .

$\Rightarrow (L_2, L_1) \in R$   
 $\therefore R$  is symmetric.

$(L_1, L_2), (L_2, L_3) \in R$   
 $\Rightarrow L_1$  is parallel to  $L_2$   
 $\Rightarrow L_2$  is parallel to  $L_3$   
 $\therefore L_1$  is parallel to  $L_3$ .  
 $\Rightarrow (L_1, L_3) \in R$   
 $\therefore R$  is transitive.

$R$  is an equivalence relation.

Set of all lines related to the line  $y = 2x + 4$  is the set of all lines that are parallel to the line  $y = 2x + 4$ .

Slope of the line  $y = 2x + 4$  is  $m = 2$ .

Line parallel to the given line is in the form  $y = 2x + c$ , where  $c \in R$ .

Set of all lines related to the given line is given by  $y = 2x + c$ , where  $c \in R$ .

### Question 15:

Let  $R$  be the relation in the set  $\{1, 2, 3, 4\}$  given by

$$R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}.$$

Choose the correct answer.

- A.  $R$  is reflexive and symmetric but not transitive.
- B.  $R$  is reflexive and transitive but not symmetric.
- C.  $R$  is symmetric and transitive but not reflexive.
- D.  $R$  is an equivalence relation.

### Solution:

$$R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$$

$$(a, a) \in R \text{ for every } a \in \{1, 2, 3, 4\}$$

$\therefore R$  is reflexive.

$(1, 2) \in R$  but  $(2, 1) \notin R$   
 $\therefore R$  is not symmetric.

$(a, b), (b, c) \in R$  for all  $a, b, c \in \{1, 2, 3, 4\}$   
 $\therefore R$  is not transitive.

$R$  is reflexive and transitive but not symmetric.

The correct answer is B.

**Question 16:**

Let  $R$  be the relation in the set  $N$  given by  $R = \{(a, b) : a = b - 2, b > 6\}$ . Choose the correct answer.

- A.  $(2, 4) \in R$
- B.  $(3, 8) \in R$
- C.  $(6, 8) \in R$
- D.  $(8, 7) \in R$

**Solution:**

$$R = \{(a, b) : a = b - 2, b > 6\}$$

Now,

$$b > 6, (2, 4) \notin R$$

$$3 \neq 8 - 2$$

$$\therefore (3, 8) \notin R \text{ and as } 8 \neq 7 - 2$$

$$\therefore (8, 7) \notin R$$

Consider  $(6, 8)$

$$8 > 6 \text{ and } 6 = 8 - 2$$

$$\therefore (6, 8) \in R$$

The correct answer is C.

## EXERCISE 1.2

### Question 1:

Show that the function  $f: R_{\bullet} \rightarrow R_{\bullet}$  defined by  $f(x) = \frac{1}{x}$  is one –one and onto, where  $R_{\bullet}$  is the set of all non –zero real numbers. Is the result true, if the domain  $R_{\bullet}$  is replaced by  $\mathbb{N}$  with co-domain being same as  $R_{\bullet}$ ?

### Solution:

$f: R_{\bullet} \rightarrow R_{\bullet}$  is by  $f(x) = \frac{1}{x}$

For one-one:

$x, y \in R_{\bullet}$  such that  $f(x) = f(y)$

$$\Rightarrow \frac{1}{x} = \frac{1}{y}$$

$$\Rightarrow x = y$$

$\therefore f$  is one-one.

For onto:

For  $y \in R$ , there exists  $x = \frac{1}{y} \in R_{\bullet}$  [as  $y \neq 0$ ] such that

$$f(x) = \frac{1}{\left(\frac{1}{y}\right)} = y$$

$\therefore f$  is onto.

Given function  $f$  is one-one and onto.

Consider function  $g: N \rightarrow R$ , defined by  $g(x) = \frac{1}{x}$

We have,  $g(x_1) = g(x_2) \Rightarrow \frac{1}{x_1} = \frac{1}{x_2} \Rightarrow x_1 = x_2$

$\therefore g$  is one-one.

$g$  is not onto as for  $1.2 \in R$ , there exist any  $x$  in  $N$  such that  $g(x) = \frac{1}{1.2}$

Function  $g$  is one-one but not onto.

### Question 2:

Check the injectivity and surjectivity of the following functions:

- i.  $f: N \rightarrow N$  given by  $f(x) = x^2$
- ii.  $f: Z \rightarrow Z$  given by  $f(x) = x^2$
- iii.  $f: R \rightarrow R$  given by  $f(x) = x^2$
- iv.  $f: N \rightarrow N$  given by  $f(x) = x^3$
- v.  $f: Z \rightarrow Z$  given by  $f(x) = x^3$

### Solution:

- i. For  $f: N \rightarrow N$  given by  $f(x) = x^2$   
 $x, y \in N$   
 $f(x) = f(y) \Rightarrow x^2 = y^2 \Rightarrow x = y$   
 $\therefore f$  is injective.

$2 \in N$ . But, there does not exist any  $x$  in  $N$  such that  $f(x) = x^2 = 2$

$\therefore f$  is not surjective

Function  $f$  is injective but not surjective.

ii.  $f : Z \rightarrow Z$  given by  $f(x) = x^2$

$$f(-1) = f(1) = 1 \text{ but } -1 \neq 1$$

$\therefore f$  is not injective.

$-2 \in Z$  But, there does not exist any  $x \in Z$  such that  $f(x) = -2 \Rightarrow x^2 = -2$

$\therefore f$  is not surjective.

Function  $f$  is neither injective nor surjective.

iii.  $f : R \rightarrow R$  given by  $f(x) = x^2$

$$f(-1) = f(1) = 1 \text{ but } -1 \neq 1$$

$\therefore f$  is not injective.

$-2 \in Z$  But, there does not exist any  $x \in Z$  such that  $f(x) = -2 \Rightarrow x^2 = -2$

$\therefore f$  is not surjective.

Function  $f$  is neither injective nor surjective.

iv.  $f : N \rightarrow N$  given by  $f(x) = x^3$

$$x, y \in N$$

$$f(x) = f(y) \Rightarrow x^3 = y^3 \Rightarrow x = y$$

$\therefore f$  is injective.

$2 \in N$ . But, there does not exist any  $x$  in  $N$  such that  $f(x) = x^3 = 2$

$\therefore f$  is not surjective

Function  $f$  is injective but not surjective.

v.  $f : Z \rightarrow Z$  given by  $f(x) = x^3$

$$x, y \in Z$$

$$f(x) = f(y) \Rightarrow x^3 = y^3 \Rightarrow x = y$$

$\therefore f$  is injective.

$2 \in Z$  But, there does not exist any  $x$  in  $Z$  such that  $f(x) = x^3 = 2$

$\therefore f$  is not surjective.

Function  $f$  is injective but not surjective.

### Question 3:

Prove that the greatest integer function  $f : R \rightarrow R$  given by  $f(x) = [x]$  is neither one-one nor onto, where  $[x]$  denotes the greatest integer less than or equal to  $x$ .

#### Solution:

$f : R \rightarrow R$  given by  $f(x) = [x]$   
 $f(1.2) = [1.2] = 1, f(1.9) = [1.9] = 1$   
 $\therefore f(1.2) = f(1.9)$ , but  $1.2 \neq 1.9$   
 $\therefore f$  is not one-one.

Consider  $0.7 \in R$

$f(x) = [x]$  is an integer. There does not exist any element  $x \in R$  such that  $f(x) = 0.7$   
 $\therefore f$  is not onto.

The greatest integer function is neither one-one nor onto.

### Question 4:

Show that the modulus function  $f : R \rightarrow R$  given by  $f(x) = |x|$  is neither one-one nor onto, where  $|x|$  is  $x$ , if  $x$  is positive or 0 and  $|x|$  is  $-x$ , if  $x$  is negative.

#### Solution:

$f : R \rightarrow R$  is  $f(x) = |x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$   
 $f(-1) = |-1| = 1$  and  $f(1) = |1| = 1$   
 $\therefore f(-1) = f(1)$  but  $-1 \neq 1$   
 $\therefore f$  is not one-one.

Consider  $-1 \in R$

$f(x) = |x|$  is non-negative. There exist any element  $x$  in domain  $R$  such that  $f(x) = |x| = -1$   
 $\therefore f$  is not onto.

The modulus function is neither one-one nor onto.

### Question 5:

$$f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$$

Show that the signum function  $f : R \rightarrow R$  given by is neither one-one nor onto.

### Solution:

$$f : R \rightarrow R \text{ is } f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$$

$$f(1) = f(2) = 1, \text{ but } 1 \neq 2$$

$\therefore f$  is not one-one.

$f(x)$  takes only 3 values  $(1, 0, -1)$  for the element  $-2$  in co-domain

$R$ , there does not exist any  $x$  in domain  $R$  such that  $f(x) = -2$ .

$\therefore f$  is not onto.

The signum function is neither one-one nor onto.

### Question 6:

Let  $A = \{1, 2, 3\}$ ,  $B = \{4, 5, 6, 7\}$  and let  $f = \{(1, 4), (2, 5), (3, 6)\}$  be a function from  $A$  to  $B$ . Show that  $f$  is one-one.

### Solution:

$$A = \{1, 2, 3\}, B = \{4, 5, 6, 7\}$$

$$f : A \rightarrow B \text{ is defined as } f = \{(1, 4), (2, 5), (3, 6)\}$$

$$\therefore f(1) = 4, f(2) = 5, f(3) = 6$$

It is seen that the images of distinct elements of  $A$  under  $f$  are distinct.

$\therefore f$  is one-one.

### Question 7:

In each of the following cases, state whether the function is one-one, onto or bijective. Justify your answer.

i.  $f : R \rightarrow R$  defined by  $f(x) = 3 - 4x$

ii.  $f : R \rightarrow R$  defined by  $f(x) = 1 + x^2$

**Solution:**

i.  $f : R \rightarrow R$  defined by  $f(x) = 3 - 4x$

$x_1, x_2 \in R$  such that  $f(x_1) = f(x_2)$

$$\Rightarrow 3 - 4x_1 = 3 - 4x_2$$

$$\Rightarrow -4x_1 = -4x_2$$

$$\Rightarrow x_1 = x_2$$

$\therefore f$  is one-one.

For any real number  $(y)$  in  $R$ , there exists  $\frac{3-y}{4}$  in  $R$  such that  $f\left(\frac{3-y}{4}\right) = 3 - 4\left(\frac{3-y}{4}\right) = y$   
 $\therefore f$  is onto.

Hence,  $f$  is bijective.

ii.  $f : R \rightarrow R$  defined by  $f(x) = 1 + x^2$

$x_1, x_2 \in R$  such that  $f(x_1) = f(x_2)$

$$\Rightarrow 1 + x_1^2 = 1 + x_2^2$$

$$\Rightarrow x_1^2 = x_2^2$$

$$\Rightarrow x_1 = \pm x_2$$

$\therefore f(x_1) = f(x_2)$  does not imply that  $x_1 = x_2$

Consider  $f(1) = f(-1) = 2$

$\therefore f$  is not one-one.

Consider an element  $-2$  in co domain  $R$ .

It is seen that  $f(x) = 1 + x^2$  is positive for all  $x \in R$ .

$\therefore f$  is not onto.

Hence,  $f$  is neither one-one nor onto.

**Question 8:**

Let  $A$  and  $B$  be sets. Show that  $f : A \times B \rightarrow B \times A$  such that  $f(a, b) = (b, a)$  is a bijective function.

**Solution:**

$f : A \times B \rightarrow B \times A$  is defined as  $f(a, b) = (b, a)$ .

$(a_1, b_1), (a_2, b_2) \in A \times B$  such that  $f(a_1, b_1) = f(a_2, b_2)$

$$\Rightarrow (b_1, a_1) = (b_2, a_2)$$

$$\Rightarrow b_1 = b_2 \text{ and } a_1 = a_2$$

$$\Rightarrow (a_1, b_1) = (a_2, b_2)$$

$\therefore f$  is one-one.

$(b, a) \in B \times A$  there exist  $(a, b) \in A \times B$  such that  $f(a, b) = (b, a)$

$\therefore f$  is onto.

$f$  is bijective.

### Question 9:

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}$$

Let  $f: N \rightarrow N$  be defined as  $f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}$  for all  $n \in N$ . State whether the function  $f$  is bijective. Justify your answer.

### Solution:

$f: N \rightarrow N$  be defined as  $f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}$  for all  $n \in N$ .

$$f(1) = \frac{1+1}{2} = 1 \text{ and } f(2) = \frac{2}{2} = 1$$

$$f(1) = f(2), \text{ where } 1 \neq 2$$

$\therefore f$  is not one-one.

Consider a natural number  $n$  in co domain  $N$ .

Case I:  $n$  is odd

$\therefore n = 2r + 1$  for some  $r \in N$  there exists  $4r + 1 \in N$  such that

$$f(4r + 1) = \frac{4r + 1 + 1}{2} = 2r + 1$$

Case II:  $n$  is even

$\therefore n = 2r$  for some  $r \in N$  there exists  $4r \in N$  such that

$$f(4r) = \frac{4r}{2} = 2r$$

$\therefore f$  is onto.

$f$  is not a bijective function.

**Question 10:**

Let  $A = \mathbb{R} - \{3\}$ ,  $B = \mathbb{R} - \{1\}$  and  $f : A \rightarrow B$  defined by  $f(x) = \left(\frac{x-2}{x-3}\right)$ . Is  $f$  one-one and onto? Justify your answer.

**Solution:**

$A = \mathbb{R} - \{3\}$ ,  $B = \mathbb{R} - \{1\}$  and  $f : A \rightarrow B$  defined by  $f(x) = \left(\frac{x-2}{x-3}\right)$

$x, y \in A$  such that  $f(x) = f(y)$

$$\Rightarrow \frac{x-2}{x-3} = \frac{y-2}{y-3}$$

$$\Rightarrow (x-2)(y-3) = (y-2)(x-3)$$

$$\Rightarrow xy - 3x - 2y + 6 = xy - 3y - 2x + 6$$

$$\Rightarrow -3x - 2y = -3y - 2x$$

$$\Rightarrow 3x - 2x = 3y - 2y$$

$$\Rightarrow x = y$$

$\therefore f$  is one-one.

Let  $y \in B = \mathbb{R} - \{1\}$ , then  $y \neq 1$

The function  $f$  is onto if there exists  $x \in A$  such that  $f(x) = y$ .

Now,

$$f(x) = y$$

$$\Rightarrow \frac{x-2}{x-3} = y$$

$$\Rightarrow x-2 = xy-3y$$

$$\Rightarrow x(1-y) = -3y+2$$

$$\Rightarrow x = \frac{2-3y}{1-y} \in A \quad [y \neq 1]$$

Thus, for any  $y \in B$ , there exists  $\frac{2-3y}{1-y} \in A$  such that

$$f\left(\frac{2-3y}{1-y}\right) = \frac{\left(\frac{2-3y}{1-y}\right)-2}{\left(\frac{2-3y}{1-y}\right)-3} = \frac{2-3y-2+2y}{2-3y-3+3y} = \frac{-y}{-1} = y$$

$\therefore f$  is onto.

Hence, the function is one-one and onto.

### Question 11:

Let  $f : R \rightarrow R$  defined as  $f(x) = x^4$ . Choose the correct answer.

- A.  $f$  is one-one onto
- B.  $f$  is many-one onto
- C.  $f$  is one-one but not onto
- D.  $f$  is neither one-one nor onto

### Solution:

$f : R \rightarrow R$  defined as  $f(x) = x^4$

$x, y \in R$  such that  $f(x) = f(y)$

$$\Rightarrow x^4 = y^4$$

$$\Rightarrow x = \pm y$$

$\therefore f(x) = f(y)$  does not imply that  $x = y$ .

For example  $f(1) = f(-1) = 1$

$\therefore f$  is not one-one.

Consider an element 2 in co domain  $R$  there does not exist any  $x$  in domain  $R$  such that  $f(x) = 2$ .

$\therefore f$  is not onto.

Function  $f$  is neither one-one nor onto.

The correct answer is D.

### Question 12:

Let  $f : R \rightarrow R$  defined as  $f(x) = 3x$ . Choose the correct answer.

- A.  $f$  is one-one onto
- B.  $f$  is many-one onto
- C.  $f$  is one-one but not onto
- D.  $f$  is neither one-one nor onto

### Solution:

$f : R \rightarrow R$  defined as  $f(x) = 3x$

$x, y \in R$  such that  $f(x) = f(y)$

$$\Rightarrow 3x = 3y$$

$$\Rightarrow x = y$$

$\therefore f$  is one-one.

For any real number  $y$  in co domain  $\mathbb{R}$ , there exist  $\frac{y}{3}$  in  $\mathbb{R}$  such that  $f\left(\frac{y}{3}\right) = 3\left(\frac{y}{3}\right) = y$

$\therefore f$  is onto.

Hence, function  $f$  is one-one and onto.

The correct answer is A.

## EXERCISE 1.3

### Question 1:

Let  $f: \{1,3,4\} \rightarrow \{1,2,5\}$  and  $g: \{1,2,5\} \rightarrow \{1,3\}$  be given by  $f = \{(1,2), (3,5), (4,1)\}$  and  $g = \{(1,3), (2,3), (5,1)\}$ . Write down  $gof$ .

### Solution:

The functions  $f: \{1,3,4\} \rightarrow \{1,2,5\}$  and  $g: \{1,2,5\} \rightarrow \{1,3\}$  are  $f = \{(1,2), (3,5), (4,1)\}$  and  $g = \{(1,3), (2,3), (5,1)\}$

$$gof(1) = g[f(1)] = g(2) = 3 \quad [as\ f(1) = 2\ and\ g(2) = 3]$$

$$gof(3) = g[f(3)] = g(5) = 1 \quad [as\ f(3) = 5\ and\ g(5) = 1]$$

$$gof(4) = g[f(4)] = g(1) = 3 \quad [as\ f(4) = 1\ and\ g(1) = 3]$$

$$\therefore gof = \{(1,3), (3,1), (4,3)\}$$

### Question 2:

Let  $f, g, h$  be functions from  $R$  to  $R$ . Show that

$$(f + g)oh = foh + goh$$

$$(f.g)oh = (foh).(goh)$$

### Solution:

$$(f + g)oh = foh + goh$$

$$LHS = [(f + g)oh](x)$$

$$= (f + g)[h(x)] = f[h(x)] + g[h(x)]$$

$$= (foh)(x) + goh(x)$$

$$= \{(foh) + (goh)\}(x) = RHS$$

$$\therefore \{(f + g)oh\}(x) = \{(foh) + (goh)\}(x) \text{ for all } x \in R$$

Hence,  $(f + g)oh = foh + goh$

$$(f.g)oh = (foh).(goh)$$

$$LHS = [(f.g)oh](x)$$

$$= (f.g)[h(x)] = f[h(x)].g[h(x)]$$

$$= (foh)(x).(goh)(x)$$

$$= \{(foh).(goh)\}(x) = RHS$$

$$\therefore [(f.g)oh](x) = \{(foh).(goh)\}(x) \text{ for all } x \in R$$

Hence,  $(f.g)oh = (foh).(goh)$

### Question 3:

Find  $gof$  and  $fog$ , if

i.  $f(x) = |x|$  and  $g(x) = |5x - 2|$

ii.  $f(x) = 8x^3$  and  $g(x) = x^{\frac{1}{3}}$

### Solution:

i.  $f(x) = |x|$  and  $g(x) = |5x - 2|$

$$\therefore gof(x) = g(f(x)) = g(|x|) = |5|x| - 2|$$

$$fog(x) = f(g(x)) = f(|5x - 2|) = ||5x - 2|| = |5x - 2|$$

ii.  $f(x) = 8x^3$  and  $g(x) = x^{\frac{1}{3}}$

$$\therefore gof(x) = g(f(x)) = g(8x^3) = (8x^3)^{\frac{1}{3}} = 2x$$

$$fog(x) = f(g(x)) = f\left(x^{\frac{1}{3}}\right)^3 = 8\left(x^{\frac{1}{3}}\right)^3 = 8x$$

### Question 4:

If  $f(x) = \frac{(4x+3)}{(6x-4)}, x \neq \frac{2}{3}$ , show that  $fof(x) = x$ , for all  $x \neq \frac{2}{3}$ . What is the reverse of  $f$ ?

### Solution:

$$(fof)(x) = f(f(x)) = f\left(\frac{4x+3}{6x-4}\right)$$

$$= \frac{4\left(\frac{4x+3}{6x-4}\right) + 3}{6\left(\frac{4x+3}{6x-4}\right) - 4} = \frac{16x+12+18x-12}{24x+18-24x+16} = \frac{34x}{34} = x$$

$$\therefore fof(x) = x \quad \text{for all } x \neq \frac{2}{3}$$

$$\Rightarrow fof = 1$$

Hence, the given function  $f$  is invertible and the inverse of  $f$  is  $f$  itself.

### Question 5:

State with reason whether the following functions have inverse.

- i.  $f: \{1, 2, 3, 4\} \rightarrow \{10\}$  with  $f = \{(1, 10), (2, 10), (3, 10), (4, 10)\}$
- ii.  $g: \{5, 6, 7, 8\} \rightarrow \{1, 2, 3, 4\}$  with  $g = \{(5, 4), (6, 3), (7, 4), (8, 2)\}$
- iii.  $h: \{2, 3, 4, 5\} \rightarrow \{7, 9, 11, 13\}$  with  $h = \{(2, 7), (3, 9), (4, 11), (5, 13)\}$

### Solution:

- i.  $f: \{1, 2, 3, 4\} \rightarrow \{10\}$  with  $f = \{(1, 10), (2, 10), (3, 10), (4, 10)\}$

$f$  is a many one function as  $f(1) = f(2) = f(3) = f(4) = 10$

$\therefore f$  is not one-one.

Function  $f$  does not have an inverse.

- ii.  $g: \{5, 6, 7, 8\} \rightarrow \{1, 2, 3, 4\}$  with  $g = \{(5, 4), (6, 3), (7, 4), (8, 2)\}$

$g$  is a many one function as  $g(5) = g(7) = 4$

$\therefore g$  is not one-one.

Function  $g$  does not have an inverse.

- iii.  $h: \{2, 3, 4, 5\} \rightarrow \{7, 9, 11, 13\}$  with  $h = \{(2, 7), (3, 9), (4, 11), (5, 13)\}$

All distinct elements of the set  $\{2, 3, 4, 5\}$  have distinct images under  $h$ .

$\therefore h$  is one-one.

$h$  is onto since for every element  $y$  of the set  $\{7, 9, 11, 13\}$ , there exists an element  $x$  in the set  $\{2, 3, 4, 5\}$ , such that  $h(x) = y$ .

$h$  is a one-one and onto function.

Function  $h$  has an inverse.

### Question 6:

Show that  $f: [-1, 1] \rightarrow R$ , given by  $f(x) = \frac{x}{(x+2)}$  is one-one. Find the inverse of the function  $f: [-1, 1] \rightarrow \text{Range } f$ .

(Hint: For  $y \in \text{Range } f$ ,  $y = f(x) = \frac{x}{x+2}$ , for some  $x$  in  $[-1, 1]$ , i.e.,  $x = \frac{2y}{(1-y)}$ )

**Solution:**

$f : [-1,1] \rightarrow R$ , given by  $f(x) = \frac{x}{(x+2)}$

For one-one

$$f(x) = f(y)$$

$$\Rightarrow \frac{x}{x+2} = \frac{y}{y+2}$$

$$\Rightarrow xy + 2x = xy + 2y$$

$$\Rightarrow 2x = 2y$$

$$\Rightarrow x = y$$

$\therefore f$  is a one-one function.

It is clear that  $f : [-1,1] \rightarrow R$  is onto.

$f : [-1,1] \rightarrow R$  is one-one and onto and therefore, the inverse of the function  $f : [-1,1] \rightarrow R$  exists.

Let  $g : \text{Range } f \rightarrow [-1,1]$  be the inverse of  $f$ .

Let  $y$  be an arbitrary element of range  $f$ .

Since  $f : [-1,1] \rightarrow \text{Range } f$  is onto, we have:

$$y = f(x) \text{ for some } x \in [-1,1]$$

$$\Rightarrow y = \frac{x}{x+2}$$

$$\Rightarrow xy + 2y = x$$

$$\Rightarrow x(1-y) = 2y$$

$$\Rightarrow x = \frac{2y}{1-y}, y \neq 1$$

Now, let us define  $g : \text{Range } f \rightarrow [-1,1]$  as

$$g(y) = \frac{2y}{1-y}, y \neq 1$$

Now,

$$(gof)(x) = g(f(x)) = g\left(\frac{x}{x+2}\right) = \frac{2\left(\frac{x}{x+2}\right)}{1 - \frac{x}{x+2}} = \frac{2x}{x+2-x} = \frac{2x}{2} = x$$

$$(fog)(x) = f(g(y)) = f\left(\frac{2y}{1-y}\right) = \frac{\frac{2y}{1-y}}{\frac{2y}{1-y} + 2} = \frac{2y}{2y+2-2y} = \frac{2y}{2} = y$$

$$\therefore gof = I_{[-1,1]} \quad \text{and} \quad fog = I_{\text{Range } f}$$

$$\therefore f^{-1} = g$$

$$\Rightarrow f^{-1}(y) = \frac{2y}{1-y}, y \neq 1$$

### Question 7:

Consider  $f: R \rightarrow R$  given by  $f(x) = 4x + 3$ . Show that  $f$  is invertible. Find the inverse of  $f$ .

### Solution:

$f: R \rightarrow R$  given by  $f(x) = 4x + 3$

For one-one

$$f(x) = f(y)$$

$$\Rightarrow 4x + 3 = 4y + 3$$

$$\Rightarrow 4x = 4y$$

$$\Rightarrow x = y$$

$\therefore f$  is a one-one function.

For onto

$$y \in R, \text{ let } y = 4x + 3$$

$$\Rightarrow x = \frac{y-3}{4} \in R$$

Therefore, for any  $y \in R$ , there exists  $x = \frac{y-3}{4} \in R$  such that

$$f(x) = f\left(\frac{y-3}{4}\right) = 4\left(\frac{y-3}{4}\right) + 3 = y$$

$\therefore f$  is onto.

Thus,  $f$  is one-one and onto and therefore,  $f^{-1}$  exists.

Let us define  $g: R \rightarrow R$  by  $g(x) = \frac{y-3}{4}$

Now,

$$(gof)(x) = g(f(x)) = g(4x+3) = \frac{(4x+3)-3}{4} = x$$

$$(fog)(y) = f(g(y)) = f\left(\frac{y-3}{4}\right) = 4\left(\frac{y-3}{4}\right) + 3 = y - 3 + 3 = y$$

$$\therefore gof = fog = I_R$$

Hence,  $f$  is invertible and the inverse of  $f$  is given by

$$f^{-1}(y) = g(y) = \frac{y-3}{4}.$$

### Question 8:

Consider  $f: R_+ \rightarrow [4, \infty)$  given by  $f(x) = x^2 + 4$ . Show that  $f$  is invertible with inverse  $f^{-1}$  of given  $f$  by  $f^{-1}(y) = \sqrt{y-4}$ , where  $R_+$  is the set of all non-negative real numbers.

### Solution:

$$f: R_+ \rightarrow [4, \infty) \text{ given by } f(x) = x^2 + 4$$

For one-one:

$$\text{Let } f(x) = f(y)$$

$$\Rightarrow x^2 + 4 = y^2 + 4$$

$$\Rightarrow x^2 = y^2$$

$$\Rightarrow x = y \quad [\text{as } x \in R]$$

$\therefore f$  is a one-one function.

For onto:

$$\text{For } y \in [4, \infty), \text{ let } y = x^2 + 4$$

$$\Rightarrow x^2 = y - 4 \geq 0 \quad [\text{as } y \geq 4]$$

$$\Rightarrow x = \sqrt{y-4} \geq 0$$

Therefore, for any  $y \in R$ , there exists  $x = \sqrt{y-4} \in R$  such that

$$f(x) = f(\sqrt{y-4}) = (\sqrt{y-4})^2 + 4 = y - 4 + 4 = y$$

$\therefore f$  is an onto function.

Thus,  $f$  is one-one and onto and therefore,  $f^{-1}$  exists.

Let us define  $g: [4, \infty) \rightarrow R_+$  by

$$g(y) = \sqrt{y-4}$$

$$\text{Now, } g \circ f(x) = g(f(x)) = g(x^2 + 4) = \sqrt{(x^2 + 4) - 4} = \sqrt{x^2} = x$$

$$\text{And } f \circ g(y) = f(g(y)) = f(\sqrt{y-4}) = (\sqrt{y-4})^2 + 4 = (y-4) + 4 = y$$

$$\therefore g \circ f = f \circ g = I_R$$

Hence,  $f$  is invertible and the inverse of  $f$  is given by

$$f^{-1}(y) = g(y) = \sqrt{y-4}.$$

### Question 9:

Consider  $f: R_+ \rightarrow [-5, \infty)$  given by  $f(x) = 9x^2 + 6x - 5$ . Show that  $f$  is invertible with

$$f^{-1}(y) = \left( \frac{(\sqrt{y+6}) - 1}{3} \right).$$

### Solution:

$$f: R_+ \rightarrow [-5, \infty) \text{ given by } f(x) = 9x^2 + 6x - 5$$

Let  $y$  be an arbitrary element of  $[-5, \infty)$ .

$$\text{Let } y = 9x^2 + 6x - 5$$

$$\Rightarrow y = (3x+1)^2 - 1 - 5$$

$$\Rightarrow y = (3x+1)^2 - 6$$

$$\Rightarrow (3x+1)^2 = y+6$$

$$\Rightarrow 3x+1 = \sqrt{y+6} \quad [\text{as } y \geq -5 \Rightarrow y+6 > 0]$$

$$\Rightarrow x = \frac{\sqrt{y+6} - 1}{3}$$

$\therefore f$  is onto, thereby range  $f = [-5, \infty)$ .

$$\text{Let us define } g: [-5, \infty) \rightarrow R_+ \text{ as } g(y) = \frac{\sqrt{y+6} - 1}{3}$$

We have,

$$\begin{aligned}
(gof)(x) &= g(f(x)) = g(9x^2 + 6x - 5) \\
&= g((3x+1)^2 - 6) \\
&= \frac{\sqrt{(3x+1)^2 - 6 + 6} - 1}{3} \\
&= \frac{3x+1-1}{3} = x
\end{aligned}$$

And,

$$\begin{aligned}
(fog)(y) &= f(g(y)) = f\left(\frac{\sqrt{y+6}-1}{3}\right) \\
&= \left[3\left(\frac{\sqrt{y+6}-1}{3}\right) + 1\right]^2 - 6 \\
&= (\sqrt{y+6})^2 - 6 = y + 6 - 6 = y
\end{aligned}$$

$$\therefore gof = I_R \text{ and } fog = I_{[-5, \infty)}$$

Hence,  $f$  is invertible and the inverse of  $f$  is given by

$$f^{-1}(y) = g(y) = \frac{\sqrt{y+6}-1}{3}.$$

### Question 10:

Let  $f : X \rightarrow Y$  be an invertible function. Show that  $f$  has unique inverse.

(Hint: suppose  $g_1$  and  $g_2$  are two inverses of  $f$ . Then for all  $y \in Y$ ,  $fog_1(y) = I_Y(y) = fog_2(y)$

. Use one-one ness of  $f$ .

### Solution:

Let  $f : X \rightarrow Y$  be an invertible function.

Also suppose  $f$  has two inverses ( $g_1$  and  $g_2$ )

Then, for all  $y \in Y$ ,

$$fog_1(y) = I_Y(y) = fog_2(y)$$

$$\Rightarrow f(g_1(y)) = f(g_2(y))$$

$$\Rightarrow g_1(y) = g_2(y) \quad [f \text{ is invertible} \Rightarrow f \text{ is one-one}]$$

$$\Rightarrow g_1 = g_2 \quad [g \text{ is one-one}]$$

Hence,  $f$  has unique inverse.

### Question 11:

Consider  $f : \{1, 2, 3\} \rightarrow \{a, b, c\}$  given by  $f(1) = a, f(2) = b, f(3) = c$ . Find  $(f^{-1})^{-1} = f$ .

### Solution:

Function  $f : \{1, 2, 3\} \rightarrow \{a, b, c\}$  given by  $f(1) = a, f(2) = b, f(3) = c$

If we define  $g : \{a, b, c\} \rightarrow \{1, 2, 3\}$  as  $g(a) = 1, g(b) = 2, g(c) = 3$

$$(f \circ g)(a) = f(g(a)) = f(1) = a$$

$$(f \circ g)(b) = f(g(b)) = f(2) = b$$

$$(f \circ g)(c) = f(g(c)) = f(3) = c$$

And,

$$(g \circ f)(1) = g(f(1)) = g(a) = 1$$

$$(g \circ f)(2) = g(f(2)) = g(b) = 2$$

$$(g \circ f)(3) = g(f(3)) = g(c) = 3$$

$$\therefore g \circ f = I_X \quad \text{and} \quad f \circ g = I_Y \quad \text{where } X = \{(1, 2, 3)\} \text{ and } Y = \{a, b, c\}$$

Thus, the inverse of  $f$  exists and  $f^{-1} = g$ .

$$\therefore f^{-1} : \{a, b, c\} \rightarrow \{1, 2, 3\} \text{ is given by, } f^{-1}(a) = 1, f^{-1}(b) = 2, f^{-1}(c) = 3$$

We need to find the inverse of  $f^{-1}$  i.e., inverse of  $g$ .

If we define  $h : \{1, 2, 3\} \rightarrow \{a, b, c\}$  as  $h(1) = a, h(2) = b, h(3) = c$

$$(g \circ h)(1) = g(h(1)) = g(a) = 1$$

$$(g \circ h)(2) = g(h(2)) = g(b) = 2$$

$$(g \circ h)(3) = g(h(3)) = g(c) = 3$$

And,

$$(h \circ g)(a) = h(g(a)) = h(1) = a$$

$$(h \circ g)(b) = h(g(b)) = h(2) = b$$

$$(h \circ g)(c) = h(g(c)) = h(3) = c$$

$$\therefore g \circ h = I_X \quad \text{and} \quad h \circ g = I_Y \quad \text{where } X = \{(1, 2, 3)\} \text{ and } Y = \{a, b, c\}$$

Thus, the inverse of  $g$  exists and  $g^{-1} = h \Rightarrow (f^{-1})^{-1} = h$ .

It can be noted that  $h = f$ .

Hence,  $(f^{-1})^{-1} = f$

### Question 12:

Let  $f : X \rightarrow Y$  be an invertible function. Show that the inverse of  $f^{-1}$  is  $f$  i.e.,  $(f^{-1})^{-1} = f$ .

### Solution:

Let  $f : X \rightarrow Y$  be an invertible function.

Then there exists a function  $g : Y \rightarrow X$  such that  $gof = I_X$  and  $fog = I_Y$

Here,  $f^{-1} = g$

Now,  $gof = I_X$  and  $fog = I_Y$

$\Rightarrow f^{-1}of = I_X$  and  $fof^{-1} = I_Y$

Hence,  $f^{-1} : Y \rightarrow X$  is invertible and  $f^{-1}$  is  $f$  i.e.,  $(f^{-1})^{-1} = f$ .

### Question 13:

If  $f : R \rightarrow R$  is given by  $f(x) = (3 - x^3)^{\frac{1}{3}}$ , then  $fof(x)$  is:

- A.  $\frac{1}{x^3}$
- B.  $x^3$
- C.  $x$
- D.  $(3 - x^3)$

### Solution:

$f : R \rightarrow R$  is given by  $f(x) = (3 - x^3)^{\frac{1}{3}}$

$f(x) = (3 - x^3)^{\frac{1}{3}}$

$$\therefore fof(x) = f(f(x)) = f\left((3 - x^3)^{\frac{1}{3}}\right) = \left[3 - \left((3 - x^3)^{\frac{1}{3}}\right)^3\right]^{\frac{1}{3}}$$

$$= \left[3 - (3 - x^3)\right]^{\frac{1}{3}} = (x^3)^{\frac{1}{3}} = x$$

$\therefore fof(x) = x$

The correct answer is C.

**Question 14:**

If  $f: R - \left\{ -\frac{4}{3} \right\} \rightarrow R$  be a function defined as  $f(x) = \frac{4x}{3x+4}$ . The inverse of  $f$  is the map  $g: \text{Range } f \rightarrow R - \left\{ -\frac{4}{3} \right\}$  given by :

A.  $g(y) = \frac{3y}{3-4y}$

B.  $g(y) = \frac{4y}{4-3y}$

C.  $g(y) = \frac{4y}{3-4y}$

D.  $g(y) = \frac{3y}{4-3y}$

**Solution:**

It is given that  $f: R - \left\{ -\frac{4}{3} \right\} \rightarrow R$  is defined as  $f(x) = \frac{4x}{3x+4}$

Let  $y$  be an arbitrary element of Range  $f$ .

Then, there exists  $x \in R - \left\{ -\frac{4}{3} \right\}$  such that  $y = f(x)$ .

$$\Rightarrow y = \frac{4x}{3x+4}$$

$$\Rightarrow 3xy + 4y = 4x$$

$$\Rightarrow x(4-3y) = 4y$$

$$\Rightarrow x = \frac{4y}{4-3y}$$

Define  $f: R - \left\{ -\frac{4}{3} \right\} \rightarrow R$  as  $g(y) = \frac{4y}{4-3y}$

Now,

$$\begin{aligned}
 (g \circ f)(x) &= g(f(x)) = g\left(\frac{4x}{3x+4}\right) \\
 &= \frac{4\left(\frac{4x}{3x+4}\right)}{4-3\left(\frac{4x}{3x+4}\right)} = \frac{16x}{12x+16-12x} \\
 &= \frac{16x}{16} = x
 \end{aligned}$$

And

$$\begin{aligned}
 (f \circ g)(x) &= (g(x)) = f\left(\frac{4y}{4-3y}\right) \\
 &= \frac{4\left(\frac{4y}{4-3y}\right)}{3\left(\frac{4y}{4-3y}\right)+4} = \frac{16y}{12y+16-12y} \\
 &= \frac{16y}{16} = y
 \end{aligned}$$

$$\therefore g \circ f = I_{R - \left\{\frac{4}{3}\right\}} \text{ and } f \circ g = I_{\text{Range } f}$$

Thus,  $g$  is the inverse of  $f$  i.e.,  $f^{-1} = g$

Hence, the inverse of  $f$  is the map  $g : \text{Range } f \rightarrow R - \left\{-\frac{4}{3}\right\}$ , which is given by  $g(y) = \frac{4y}{4-3y}$ .

The correct answer is B.

## EXERCISE 1.4

### Question 1:

Determine whether or not each of the definition of  $*$  given below gives a binary operation. In the event that  $*$  is not a binary operation, give justification for this.

- i. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = a - b$
- ii. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = ab$
- iii. On  $\mathbf{R}$ , define  $*$  by  $a * b = ab^2$
- iv. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = |a - b|$
- v. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = a$

### Solution:

- i. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = a - b$

It is not a binary operation as the image of  $(1, 2)$  under  $*$  is

$$1 * 2 = 1 - 2$$

$$\Rightarrow -1 \notin \mathbf{Z}^+.$$

Therefore,  $*$  is not a binary operation.

- ii. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = ab$

It is seen that for each  $a, b \in \mathbf{Z}^+$ , there is a unique element  $ab$  in  $\mathbf{Z}^+$ .

This means that  $*$  carries each pair  $(a, b)$  to a unique element  $a * b = ab$  in  $\mathbf{Z}^+$ .

Therefore,  $*$  is a binary operation.

- iii. On  $\mathbf{R}$ , define  $*$   $a * b = ab^2$

It is seen that for each  $a, b \in \mathbf{R}$ , there is a unique element  $ab^2$  in  $\mathbf{R}$ . This means that  $*$

carries each pair  $(a, b)$  to a unique element  $a * b = ab^2$  in  $\mathbf{R}$ .

Therefore,  $*$  is a binary operation.

- iv. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = |a - b|$

It is seen that for each  $a, b \in \mathbf{Z}^+$ , there is a unique element  $|a - b|$  in  $\mathbf{Z}^+$ . This means that

$*$  carries each pair  $(a, b)$  to a unique element  $a * b = |a - b|$  in  $\mathbf{Z}^+$ . Therefore,  $*$  is a binary operation.

- v. On  $\mathbf{Z}^+$ , define  $*$  by  $a * b = a$

$*$  carries each pair  $(a, b)$  to a unique element in  $a * b = a$  in  $\mathbf{Z}^+$ .

Therefore,  $*$  is a binary operation.

### Question 2:

For each binary operation  $*$  defined below, determine whether  $*$  is commutative or associative.

- i. On  $\mathbf{Z}^+$ , define  $a * b = a - b$

- ii. On  $\mathbf{Q}$ , define  $a * b = ab + 1$
- iii. On  $\mathbf{Q}$ , define  $a * b = \frac{ab}{2}$
- iv. On  $\mathbf{Z}^+$ , define  $a * b = 2^{ab}$
- v. On  $\mathbf{Z}^+$ , define  $a * b = a^b$
- vi. On  $\mathbf{R} - \{-1\}$ , define  $a * b = \frac{a}{b+1}$

**Solution:**

- i. On  $\mathbf{Z}^+$ , define  $a * b = a - b$

It can be observed that  $1 * 2 = 1 - 2 = -1$  and  $2 * 1 = 2 - 1 = 1$ .

$\therefore 1 * 2 \neq 2 * 1$ ; where  $1, 2 \in \mathbf{Z}$

Hence, the operation  $*$  is not commutative.

Also,

$$(1 * 2) * 3 = (1 - 2) * 3 = -1 * 3 = -1 - 3 = -4$$

$$1 * (2 * 3) = 1 * (2 - 3) = 1 * -1 = 1 - (-1) = 2$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in \mathbf{Z}$

Hence, the operation  $*$  is not associative.

- ii. On  $\mathbf{Q}$ , define  $a * b = ab + 1$

$$ab = ba \quad \text{for all } a, b \in \mathbf{Q}$$

$$\Rightarrow ab + 1 = ba + 1 \quad \text{for all } a, b \in \mathbf{Q}$$

$$\Rightarrow a * b = b * a \quad \text{for all } a, b \in \mathbf{Q}$$

Hence, the operation  $*$  is commutative.

$$(1 * 2) * 3 = (1 \times 2 + 1) * 3 = 3 * 3 = 3 \times 3 + 1 = 10$$

$$1 * (2 * 3) = 1 * (2 \times 3 + 1) = 1 * 7 = 1 \times 7 + 1 = 8$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in \mathbf{Q}$

Hence, the operation  $*$  is not associative.

- iii. On  $\mathbf{Q}$ , define  $a * b = \frac{ab}{2}$

$$ab = ba \quad \text{for all } a, b \in \mathbf{Q}$$

$$\Rightarrow \frac{ab}{2} = \frac{ab}{2} \quad \text{for all } a, b \in \mathbf{Q}$$

$$\Rightarrow a * b = b * a \quad \text{for all } a, b \in \mathbf{Q}$$

Hence, the operation  $*$  is commutative.

$$(a * b) * c = \left(\frac{ab}{2}\right) * c = \frac{\left(\frac{ab}{2}\right)^c}{2} = \frac{abc}{4}$$

And

$$a * (b * c) = a * \left(\frac{bc}{2}\right) = \frac{a\left(\frac{bc}{2}\right)}{2} = \frac{abc}{4}$$

$$\therefore (a * b) * c = a * (b * c)$$

where  $a, b, c \in \mathbf{Q}$

Hence, the operation  $*$  is associative.

iv. On  $\mathbf{Z}^+$ , define  $a * b = 2^{ab}$

$$ab = ba \quad \text{for all } a, b \in \mathbf{Z}$$

$$\Rightarrow 2^{ab} = 2^{ba} \quad \text{for all } a, b \in \mathbf{Z}$$

$$\Rightarrow a * b = b * a \quad \text{for all } a, b \in \mathbf{Z}$$

Hence, the operation  $*$  is commutative.

$$(1 * 2) * 3 = 2^{1 \times 2} * 3 = 4 * 3 = 2^{4 \times 3} = 2^{12}$$

$$1 * (2 * 3) = 1 * 2^{2 \times 3} = 1 * 2^6 = 1 * 64 = 2^{64}$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in \mathbf{Z}^+$

Hence, the operation  $*$  is not associative.

v. On  $\mathbf{Z}^+$ , define  $a * b = a^b$

$$1 * 2 = 1^2 = 1$$

$$2 * 1 = 2^1 = 2$$

$$\therefore 1 * 2 \neq 2 * 1$$

where  $1, 2, \in \mathbf{Z}^+$

Hence, the operation  $*$  is not commutative.

$$(2 * 3) * 4 = 2^3 * 4 = 8 * 4 = 8^4 = 2^{12}$$

$$2 * (3 * 4) = 2 * 3^4 = 2 * 81 = 2^{81}$$

$$\therefore (2 * 3) * 4 \neq 2 * (3 * 4)$$

where  $2, 3, 4 \in \mathbf{Z}^+$

Hence, the operation  $*$  is not associative.

vi. On  $\mathbf{R} - \{-1\}$ , define  $a * b = \frac{a}{b+1}$

$$1 * 2 = \frac{1}{2+1} = \frac{1}{3}$$

$$2 * 1 = \frac{2}{1+1} = \frac{2}{2} = 1$$

$$\therefore 1 * 2 \neq 2 * 1$$

where  $1, 2 \in \mathbf{R} - \{-1\}$

Hence, the operation  $*$  is not commutative.

$$(1 * 2) * 3 = \frac{1}{3} * 3 = \frac{\frac{1}{3}}{\frac{1}{3} + 1} = \frac{1}{12}$$

$$1 * (2 * 3) = 1 * \frac{2}{3+1} = 1 * \frac{2}{4} = 1 * \frac{1}{2} = \frac{1}{\frac{1}{2} + 1} = \frac{1}{\frac{3}{2}} = \frac{2}{3}$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in \mathbf{R} - \{-1\}$

Hence, the operation  $*$  is not associative.

### Question 3:

Consider the binary operation  $\wedge$  on the set  $\{1, 2, 3, 4, 5\}$  defined by  $a \wedge b = \min\{a, b\}$ . Write the operation table of the operation  $\wedge$ .

**Solution:**

The binary operation  $\wedge$  on the set  $\{1, 2, 3, 4, 5\}$  is defined by  $a \wedge b = \min\{a, b\}$  for all  $a, b \in \{1, 2, 3, 4, 5\}$ .

The operation table for the given operation  $\wedge$  can be given as:

$\wedge$	1	2	3	4	5
1	1	1	1	1	1
2	1	2	2	2	2
3	1	2	3	3	3
4	1	2	3	4	4
5	1	2	3	4	5

### Question 4:

Consider a binary operation  $*$  on the set  $\{1, 2, 3, 4, 5\}$  given by the following multiplication table.

- Compute  $(2 * 3) * 4$  and  $2 * (3 * 4)$
- Is  $*$  commutative?
- Compute  $(2 * 3) * (4 * 5)$ .  
(Hint: Use the following table)

$*$	1	2	3	4	5
1	1	1	1	1	1
2	1	2	1	2	1

3	1	1	3	1	1
4	1	2	1	4	1
5	1	1	1	1	5

**Solution:**

$$(2 * 3) * 4 = 1 * 4 = 1$$

- i.  $2 * (3 * 4) = 2 * 1 = 1$
- ii. For every  $a, b \in \{1, 2, 3, 4, 5\}$ , we have  $a * b = b * a$ . Therefore,  $*$  is commutative.
- iii.  $(2 * 3) * (4 * 5)$   
 $(2 * 3) = 1$  and  $(4 * 5) = 1$   
 $\therefore (2 * 3) * (4 * 5) = 1 * 1 = 1$

**Question 5:**

Let  $*$ ' be the binary operation on the set  $\{1, 2, 3, 4, 5\}$  defined by  $a *' b = \text{H.C.F. of } a \text{ and } b$ . Is the operation  $*$ ' same as the operation  $*$  defined in Exercise 4 above? Justify your answer.

**Solution:**

The binary operation on the set  $\{1, 2, 3, 4, 5\}$  is defined by  $a *' b = \text{H.C.F. of } a \text{ and } b$ . The operation table for the operation  $*$ ' can be given as:

$*$ '	1	2	3	4	5
1	1	1	1	1	1
2	1	2	1	2	1
3	1	1	3	1	1
4	1	2	1	4	1
5	1	1	1	1	5

The operation table for the operations  $*$ ' and  $*$  are same. operation  $*$ ' is same as operation  $*$ .

**Question 6:**

Let  $*$  be the binary operation on  $\mathbb{N}$  defined by  $a * b = \text{L.C.M. of } a \text{ and } b$ . Find

- i.  $5 * 7, 20 * 16$
- ii. Is  $*$  commutative?
- iii. Is  $*$  associative?
- iv. Find the identity of  $*$  in  $\mathbb{N}$
- v. Which elements of  $\mathbb{N}$  are invertible for the operation  $*$ ?

**Solution:**

The binary operation on N is defined by  $a * b = \text{L.C.M. of } a \text{ and } b$ .

i.  $5 * 7 = \text{L.C.M of } 5 \text{ and } 7 = 35$

$20 * 16 = \text{LCM of } 20 \text{ and } 16 = 80$

ii.  $\text{L.C.M. of } a \text{ and } b = \text{LCM of } b \text{ and } a \text{ for all } a, b \in N$

$\therefore a * b = b * a$

Operation  $*$  is commutative.

iii. For  $a, b, c \in N$

$(a * b) * c = (\text{L.C.M. of } a \text{ and } b) * c = \text{L.C.M. of } a, b, c$

$a * (b * c) = a * (\text{L.C.M. of } b \text{ and } c) = \text{L.C.M. of } a, b, c$

$\therefore (a * b) * c = a * (b * c)$

Operation  $*$  is associative.

iv.  $\text{L.C.M. of } a \text{ and } 1 = a = \text{L.C.M. of } 1 \text{ and } a \text{ for all } a \in N$

$a * 1 = a = 1 * a \text{ for all } a \in N$

Therefore, 1 is the identity of  $*$  in N.

v. An element  $a$  in N is invertible with respect to the operation  $*$  if there exists an element  $b$  in N, such that  $a * b = e = b * a$

$e = 1$

$\text{L.C.M. of } a \text{ and } b = 1 = \text{LCM of } b \text{ and } a \text{ possible only when } a \text{ and } b \text{ are equal to } 1.$

1 is the only invertible element of N with respect to the operation  $*$ .

**Question 7:**

Is  $*$  defined on the set  $\{1, 2, 3, 4, 5\}$  by  $a * b = \text{LCM of } a \text{ and } b$  a binary operation? Justify your answer.

**Solution:**

The operation  $*$  on the set  $\{1, 2, 3, 4, 5\}$  is defined by  $a * b = \text{LCM of } a \text{ and } b$ .

The operation table for the operation  $*$  can be given as:

*	1	2	3	4	5
1	1	2	3	4	5
2	2	2	6	4	10
3	3	6	3	12	15
4	4	4	12	4	20
5	5	10	15	20	5

$$3 * 2 = 2 * 3 = 6 \notin A,$$

$$5 * 2 = 2 * 5 = 10 \notin A,$$

$$3 * 4 = 4 * 3 = 12 \notin A,$$

$$3 * 5 = 5 * 3 = 15 \notin A,$$

$$4 * 5 = 5 * 4 = 20 \notin A$$

The given operation  $*$  is not a binary operation.

### Question 8:

Let  $*$  be the binary operation on  $N$  defined by  $a * b = \text{H.C.F. of } a \text{ and } b$ . Is  $*$  commutative? Is  $*$  associative? Does there exist identity for this binary operation on  $N$ ?

### Solution:

The binary operation  $*$  on  $N$  defined by  $a * b = \text{H.C.F. of } a \text{ and } b$ .

$$\therefore a * b = b * a$$

Operation  $*$  is commutative.

For all  $a, b, c \in N$ ,

$$(a * b) * c = (\text{HCF of } a \text{ and } b) * c = \text{HCF of } a, b, c$$

$$a * (b * c) = a * (\text{HCF. of } b \text{ and } c) = \text{HCF of } a, b, c$$

$$\therefore (a * b) * c = a * (b * c)$$

Operation  $*$  is associative.

$e \in N$  will be the identity for the operation  $*$  if  $a * e = a = e * a$  for all  $a \in N$ . But this relation is not true for any  $a \in N$ .

Operation  $*$  does not have any identity in  $N$ .

### Question 9:

Let  $*$  be the binary operation on  $Q$  of rational numbers as follows:

i.  $a * b = a - b$

ii.  $a * b = a^2 + b^2$

iii.  $a * b = a + ab$

iv.  $a * b = (a - b)^2$

v.  $a + b = \frac{ab}{4}$

vi.  $a * b = ab^2$

Find which of the binary operations are commutative and which are associative.

**Solution:**

i. On  $Q$ , the operation  $*$  is defined as  $a * b = a - b$

$$\frac{1}{2} * \frac{1}{3} = \frac{1}{2} - \frac{1}{3} = \frac{3-2}{3} = \frac{1}{3}$$

And

$$\frac{1}{3} * \frac{1}{2} = \frac{1}{3} - \frac{1}{2} = \frac{2-3}{6} = \frac{-1}{6}$$

$$\therefore \left( \frac{1}{2} * \frac{1}{3} \right) \neq \left( \frac{1}{3} * \frac{1}{2} \right)$$

where  $\frac{1}{2}, \frac{1}{3} \in Q$

Operation  $*$  is not commutative.

$$\left( \frac{1}{2} * \frac{1}{3} \right) * \frac{1}{4} = \left( \frac{1}{2} - \frac{1}{3} \right) * \frac{1}{4} = \frac{1}{6} * \frac{1}{4} = \frac{1}{6} - \frac{1}{4} = \frac{2-3}{12} = \frac{-1}{12}$$

$$\frac{1}{2} * \left( \frac{1}{3} * \frac{1}{4} \right) = \frac{1}{2} * \left( \frac{1}{3} - \frac{1}{4} \right) = \frac{1}{2} * \frac{1}{12} = \frac{1}{2} - \frac{1}{12} = \frac{6-1}{12} = \frac{5}{12}$$

$$\therefore \left( \frac{1}{2} * \frac{1}{3} \right) * \frac{1}{4} \neq \frac{1}{2} * \left( \frac{1}{3} * \frac{1}{4} \right)$$

where  $\frac{1}{2}, \frac{1}{3}, \frac{1}{4} \in Q$

Operation  $*$  is not associative.

ii. On  $Q$ , the operation  $*$  is defined as  $a * b = a^2 + b^2$

For  $a, b \in Q$

$$a * b = a^2 + b^2 = b^2 + a^2 = b * a$$

$$\therefore a * b = b * a$$

Operation  $*$  is commutative.

$$(1 * 2) * 3 = (1^2 + 2^2) * 3 = (1 + 4) * 3 = 5 * 3 = 5^2 + 3^2 = 25 + 9 = 34$$

$$1 * (2 * 3) = 1 * (2^2 + 3^2) = 1 * (4 + 9) = 1 * 13 = 1^2 + 13^2 = 1 + 169 = 170$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in Q$

Operation  $*$  is not associative.

iii. On  $Q$ , the operation  $*$  is defined as  $a * b = a + ab$

$$1 * 2 = 1 + 1 \times 2 = 1 + 2 = 3$$

$$2 * 1 = 2 + 2 \times 1 = 2 + 2 = 4$$

$$\therefore 1 * 2 \neq 2 * 1$$

where  $1, 2 \in Q$

Operation  $*$  is not commutative.

$$(1 * 2) * 3 = (1 + 1 \times 2) * 3 = 3 * 3 = 3 + 3 \times 3 = 3 + 9 = 12$$

$$1 * (2 * 3) = 1 * (2 + 2 \times 3) = 1 * 8 = 1 + 1 \times 8 = 1 + 8 = 9$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in Q$

Operation  $*$  is not associative.

iv. On  $Q$ , the operation  $*$  is defined as  $a * b = (a - b)^2$

For  $a, b \in Q$

$$a * b = (a - b)^2$$

$$b * a = (b - a)^2 = [-(a - b)]^2 = (a - b)^2$$

$$\therefore a * b = b * a$$

Operation  $*$  is commutative.

$$(1 * 2) * 3 = (1 - 2)^2 * 3 = (-1)^2 * 3 = 1 * 3 = (1 - 3)^2 = (-2)^2 = 4$$

$$1 * (2 * 3) = 1 * (2 - 3)^2 = 1 * (-1)^2 = 1 * 1 = (1 - 1)^2 = 0$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

where  $1, 2, 3 \in Q$

Operation  $*$  is not associative.

v. On  $Q$ , the operation  $*$  is defined as  $a * b = \frac{ab}{4}$

For  $a, b \in Q$

$$a * b = \frac{ab}{4} = \frac{ba}{4} = b * a$$

$$\therefore a * b = b * a$$

Operation  $*$  is commutative.

For  $a, b, c \in Q$

$$(a * b) * c = \frac{ab}{4} * c = \frac{\frac{ab}{4} \cdot c}{4} = \frac{abc}{16}$$

$$a * (b * c) = a * \frac{bc}{4} = \frac{a \cdot \frac{bc}{4}}{4} = \frac{abc}{16}$$

$$\therefore (a * b) * c = a * (b * c)$$

where  $a, b, c \in Q$

Operation  $*$  is associative.

vi. On  $Q$ , the operation  $*$  is defined as  $a * b = ab^2$

$$\frac{1}{2} * \frac{1}{3} = \frac{1}{2} \cdot \left(\frac{1}{3}\right)^2 = \frac{1}{2} \cdot \frac{1}{9} = \frac{1}{18}$$

$$\frac{1}{3} * \frac{1}{2} = \frac{1}{3} \cdot \left(\frac{1}{2}\right)^2 = \frac{1}{3} \cdot \frac{1}{4} = \frac{1}{12}$$

$$\therefore \left(\frac{1}{2} * \frac{1}{3}\right) \neq \left(\frac{1}{3} * \frac{1}{2}\right)$$

where  $\frac{1}{2}, \frac{1}{3} \in Q$

Operation  $*$  is not commutative.

$$\left(\frac{1}{2} * \frac{1}{3}\right) * \frac{1}{4} = \left(\frac{1}{2} \cdot \left(\frac{1}{3}\right)^2\right) * \frac{1}{4} = \frac{1}{18} * \frac{1}{4} = \frac{1}{18} \cdot \left(\frac{1}{4}\right)^2 = \frac{1}{18 \times 16}$$

$$\frac{1}{2} * \left(\frac{1}{3} * \frac{1}{4}\right) = \frac{1}{2} * \left(\frac{1}{3} \cdot \left(\frac{1}{4}\right)^2\right) = \frac{1}{2} * \frac{1}{48} = \frac{1}{2} \cdot \left(\frac{1}{48}\right)^2 = \frac{1}{2 \times (48)^2}$$

$$\therefore \left(\frac{1}{2} * \frac{1}{3}\right) * \frac{1}{4} \neq \frac{1}{2} * \left(\frac{1}{3} * \frac{1}{4}\right) \quad \text{where } \frac{1}{2}, \frac{1}{3}, \frac{1}{4} \in Q$$

Operation  $*$  is not associative.

Operations defined in (ii), (iv), (v) are commutative and the operation defined in (v) is associative.

### Question 10:

Find which of the operations given above has identity.

### Solution:

An element  $e \in Q$  will be the identity element for the operation  $*$  if

$$a * e = a = e * a, \text{ for all } a \in Q$$

$$a * b = \frac{ab}{4}$$

$$\Rightarrow a * e = a$$

$$\Rightarrow \frac{ae}{4} = a$$

$$\Rightarrow e = 4$$

Similarly, it can be checked for  $e * a = a$ , we get  $e = 4$  is the identity.

### Question 11:

$A = N \times N$  and  $*$  be the binary operation on  $A$  defined by  $(a, b) * (c, d) = (a + c, b + d)$ . Show that  $*$  is commutative and associative. Find the identity element for  $*$  on  $A$ , if any.

### Solution:

$A = N \times N$  and  $*$  be the binary operation on  $A$  defined by

$$(a,b)*(c,d)=(a+c,b+d)$$

$$(a,b)*(c,d) \in A$$

$$a,b,c,d \in N$$

$$(a,b)*(c,d)=(a+c,b+d)$$

$$(c,d)*(a,b)=(c+a,d+b)=(a+c,b+d)$$

$$\therefore (a,b)*(c,d)=(c,d)*(a,b)$$

Operation  $*$  is commutative.

Now, let  $(a,b), (c,d), (e,f) \in A$

$a,b,c,d,e,f \in N$

$$[(a,b) * (c,d)] * (e,f) = (a+c, b+d) * (e,f) = (a+c+e, b+d+f)$$

$$(a,b) * [(c,d) * (e,f)] = (a,b) * (c+e, d+f) = (a+c+e, b+d+f)$$

$$\therefore [(a,b) * (c,d)] * (e,f) = (a,b) * [(c,d) * (e,f)]$$

Operation  $*$  is associative.

An element  $e = (e_1, e_2) \in A$  will be an identity element for the operation  $*$  if  $a + e = a = e * a$  for all  $a = (a_1, a_2) \in A$  i.e.,  $(a_1 + e_1, a_2 + e_2) = (a_1, a_2) = (e_1 + a_1, e_2 + a_2)$ , which is not true for any element in  $A$ .

Therefore, the operation  $*$  does not have any identity element.

### Question 12:

State whether the following statements are true or false. Justify.

- For an arbitrary binary operation  $*$  on a set  $N$ ,  $a * a = a$  for all  $a \in N$ .
- If  $*$  is a commutative binary operation on  $N$ , then  $a * (b * c) = (c * b) * a$

### Solution:

- Define operation  $*$  on a set  $N$  as  $a * a = a$  for all  $a \in N$ .

In particular, for  $a = 3$ ,

$$3 * 3 = 9 \neq 3$$

Therefore, statement (i) is false.

- R.H.S. =  $(c * b) * a$   
 $= (b * c) * a$  [ $*$  is commutative]  
 $= a * (b * c)$  [Again, as  $*$  is commutative]  
= L.H.S.

$$\therefore a * (b * c) = (c * b) * a$$

Therefore, statement (ii) is true.

### Question 13:

Consider a binary operation  $*$  on  $N$  defined as  $a * b = a^3 + b^3$ . Choose the correct answer.

- Is  $*$  both associative and commutative?
- Is  $*$  commutative but not associative?
- Is  $*$  associative but not commutative?
- Is  $*$  neither commutative nor associative?

**Solution:**

On  $\mathbb{N}$ , operation  $*$  is defined as  $a * b = a^3 + b^3$ .

For all  $a, b \in \mathbb{N}$

$$a * b = a^3 + b^3 = b^3 + a^3 = b * a$$

Operation  $*$  is commutative.

$$(1 * 2) * 3 = (1^3 + 2^3) * 3 = (1 + 8) * 3 = 9 * 3 = 9^3 + 3^3 = 729 + 27 = 756$$

$$1 * (2 * 3) = 1 * (2^3 + 3^3) = 1 * (8 + 27) = 1 * 35 = 1^3 + 35^3 = 1 + 42875 = 42876$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3)$$

associative.

Operation  $*$  is not

Therefore, Operation  $*$  is commutative, but not associative.

The correct answer is B.

## MISCELLANEOUS EXERCISE

### Question 1:

Let  $f: R \rightarrow R$  be defined as  $f(x) = 10x + 7$ . Find the function  $g: R \rightarrow R$  such that  $gof = fof = I_R$ .

### Solution:

$f: R \rightarrow R$  is defined as  $f(x) = 10x + 7$

For one-one:

$$f(x) = f(y) \text{ where } x, y \in R$$

$$\Rightarrow 10x + 7 = 10y + 7$$

$$\Rightarrow x = y$$

$\therefore f$  is one-one.

For onto:

$$y \in R, \text{ Let } y = 10x + 7$$

$$\Rightarrow x = \frac{y-7}{10} \in R$$

For any  $y \in R$ , there exists  $x = \frac{y-7}{10} \in R$  such that

$$f(x) = f\left(\frac{y-7}{10}\right) = 10\left(\frac{y-7}{10}\right) + 7 = y - 7 + 7 = y$$

$\therefore f$  is onto.

Thus,  $f$  is an invertible function.

Let us define  $g: R \rightarrow R$  as  $g(y) = \frac{y-7}{10}$ .

Now,

$$gof(x) = g(f(x)) = g(10x + 7) = \frac{(10x + 7) - 7}{10} = \frac{10x}{10} = x$$

And,

$$fog(y) = f(g(y)) = f\left(\frac{y-7}{10}\right) = 10\left(\frac{y-7}{10}\right) + 7 = y - 7 + 7 = y$$

$\therefore gof = I_R$  and  $fof = I_R$

Hence, the required function  $g: R \rightarrow R$  as  $g(y) = \frac{y-7}{10}$ .

## Question 2:

Let  $f : W \rightarrow W$  be defined as  $f(n) = n - 1$ , if  $n$  is odd and  $f(n) = n + 1$ , if  $n$  is even. Show that  $f$  is invertible. Find the inverse of  $f$ . Here,  $W$  is the set of all whole numbers.

### Solution:

$f : W \rightarrow W$  is defined as  $f(n) = \begin{cases} n - 1, & \text{If } n \text{ is odd} \\ n + 1, & \text{If } n \text{ is even} \end{cases}$

For one-one:

$$f(n) = f(m)$$

If  $n$  is odd and  $m$  is even, then we will have  $n - 1 = m + 1$ .

$$\Rightarrow n - m = 2$$

Similarly, the possibility of  $n$  being even and  $m$  being odd can also be ignored under a similar argument.

$\therefore$  Both  $n$  and  $m$  must be either odd or even.

Now, if both  $n$  and  $m$  are odd, then we have:

$$f(n) = f(m)$$

$$\Rightarrow n - 1 = m - 1$$

$$\Rightarrow n = m$$

Again, if both  $n$  and  $m$  are even, then we have:

$$f(n) = f(m)$$

$$\Rightarrow n + 1 = m + 1$$

$$\Rightarrow n = m$$

$\therefore f$  is one-one.

For onto:

Any odd number  $2r + 1$  in co-domain  $N$  is the image of  $2r$  in domain  $N$  and any even number  $2r$  in co-domain  $N$  is the image of  $2r + 1$  in domain  $N$ .

$\therefore f$  is onto.

$f$  is an invertible function.

Let us define  $g : W \rightarrow W$  as  $f(m) = \begin{cases} m - 1, & \text{If } m \text{ is odd} \\ m + 1, & \text{If } m \text{ is even} \end{cases}$

When  $r$  is odd

$$g \circ f(n) = g(f(n)) = g(n - 1) = n - 1 + 1 = n$$

When  $r$  is even

$$gof(n) = g(f(n)) = g(n+1) = n+1-1 = n$$

When  $m$  is odd

$$fog(m) = f(g(m)) = f(m-1) = m-1+1 = m$$

When  $m$  is even

$$fog(m) = f(g(m)) = f(m+1) = m+1-1 = m$$

$$\therefore gof = I_W \text{ and } fog = I_W$$

$f$  is invertible and the inverse of  $f$  is given by  $f^{-1} = g$ , which is the same as  $f$ .  
inverse of  $f$  is  $f$  itself.

### Question 3:

If  $f: R \rightarrow R$  be defined as  $f(x) = x^2 - 3x + 2$ , find  $f(f(x))$ .

### Solution:

$f: R \rightarrow R$  is defined as  $f(x) = x^2 - 3x + 2$ .

$$\begin{aligned} f(f(x)) &= f(x^2 - 3x + 2) \\ &= (x^2 - 3x + 2)^2 - 3(x^2 - 3x + 2) + 2 \\ &= (x^4 + 9x^2 + 4 - 6x^3 - 12x + 4x^2) + (-3x^2 + 9x - 6) + 2 \\ &= x^4 - 6x^3 + 10x^2 - 3x \end{aligned}$$

### Question 4:

Show that function  $f: R \rightarrow \{x \in R: -1 < x < 1\}$  be defined by  $f(x) = \frac{x}{1+|x|}$ ,  $x \in R$  is one-one and onto function.

### Solution:

$f: R \rightarrow \{x \in R: -1 < x < 1\}$  is defined by  $f(x) = \frac{x}{1+|x|}$ ,  $x \in R$ .

For one-one:

$$f(x) = f(y) \quad \text{where } x, y \in R$$

$$\Rightarrow \frac{x}{1+|x|} = \frac{y}{1+|y|}$$

If  $x$  is positive and  $y$  is negative,

$$\frac{x}{1+|x|} = \frac{y}{1+|y|}$$

$$\Rightarrow 2xy = x - y$$

Since,  $x$  is positive and  $y$  is negative,

$$x > y \Rightarrow x - y > 0$$

$2xy$  is negative.

$$2xy \neq x - y$$

Case of  $x$  being positive and  $y$  being negative, can be ruled out.

$\therefore x$  and  $y$  have to be either positive or negative.

If  $x$  and  $y$  are positive,

$$f(x) = f(y)$$

$$\Rightarrow \frac{x}{1+x} = \frac{y}{1+y}$$

$$\Rightarrow x - xy = y - xy$$

$$\Rightarrow x = y$$

$\therefore f$  is one-one.

For onto:

Let  $y \in R$  such that  $-1 < y < 1$ .

If  $x$  is negative, then there exists  $x = \frac{y}{1+y} \in R$  such that

$$f(x) = f\left(\frac{y}{1+y}\right) = \frac{\left(\frac{y}{1+y}\right)}{1+\left|\frac{y}{1+y}\right|} = \frac{\frac{y}{1+y}}{1+\left(\frac{-y}{1+y}\right)} = \frac{y}{1+y-y} = y$$

If  $x$  is positive, then there exists  $x = \frac{y}{1-y} \in R$  such that

$$f(x) = f\left(\frac{y}{1-y}\right) = \frac{\left(\frac{y}{1-y}\right)}{1 + \left|\frac{y}{1-y}\right|} = \frac{\frac{y}{1-y}}{1 + \left(\frac{y}{1-y}\right)} = \frac{y}{1-y+y} = y$$

∴  $f$  is onto.

Hence,  $f$  is one-one and onto.

### Question 5:

Show that function  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $f(x) = x^3$  is injective.

#### Solution:

$f: \mathbb{R} \rightarrow \mathbb{R}$  is defined by  $f(x) = x^3$

For one-one:

$$f(x) = f(y) \quad \text{where } x, y \in \mathbb{R}$$

$$x^3 = y^3 \dots\dots\dots(1)$$

We need to show that  $x = y$

Suppose  $x \neq y$ , their cubes will also not be equal.

$$\Rightarrow x^3 \neq y^3$$

This will be a contradiction to (1).

∴  $x = y$ . Hence,  $f$  is injective.

### Question 6:

Give examples of two functions  $f: \mathbb{N} \rightarrow \mathbb{Z}$  and  $g: \mathbb{Z} \rightarrow \mathbb{Z}$  such that  $g \circ f$  is injective but  $g$  is not injective.

(Hint: Consider  $f(x) = x$  and  $g(x) = |x|$ )

#### Solution:

Define  $f: \mathbb{N} \rightarrow \mathbb{Z}$  as  $f(x) = x$  and  $g: \mathbb{Z} \rightarrow \mathbb{Z}$  as  $g(x) = |x|$

Let us first show that  $g$  is not injective.

$$(-1) = |-1| = 1$$

$$(1) = |1| = 1$$

$$\therefore (-1) = g(1), \text{ but } -1 \neq 1$$

$\therefore g$  is not injective.

$g \circ f : N \rightarrow Z$  is defined as  $g \circ f(x) = g(f(x)) = g(x) = |x|$

$x, y \in N$  such that  $g \circ f(x) = g \circ f(y)$

$$\Rightarrow |x| = |y|$$

Since  $x, y \in N$ , both are positive.

$$\therefore |x| = |y|$$

$$\Rightarrow x = y$$

$\therefore g \circ f$  is injective.

### Question 7:

Given examples of two functions  $f : N \rightarrow N$  and  $g : N \rightarrow N$  such that  $g \circ f$  is onto but  $f$  is not onto.

(Hint: Consider  $f(x) = x+1$  and  $g(x) = \begin{cases} x-1, & \text{if } x > 1 \\ 1, & \text{if } x = 1 \end{cases}$ )

**Solution:**

Define  $f : N \rightarrow Z$  as  $f(x) = x+1$  and  $g : Z \rightarrow Z$  as  $g(x) = \begin{cases} x-1, & \text{if } x > 1 \\ 1, & \text{if } x = 1 \end{cases}$

Let us first show that  $g$  is not onto.

Consider element 1 in co-domain  $N$ . This element is not an image of any of the elements in domain  $N$ .

$\therefore f$  is not onto.

$g : N \rightarrow N$  is defined by

$$g \circ f(x) = g(f(x)) = g(x+1) = x+1-1 = x \quad [x \in N \Rightarrow x+1 > 1]$$

For  $y \in N$ , there exists  $x = y \in N$  such that  $g \circ f(x) = y$ .

$\therefore g \circ f$  is onto.

### Question 8:

Given a non-empty set  $X$ , consider  $P(X)$  which is the set of all subsets of  $X$ .

Define the relation  $R$  in  $P(X)$  as follows:

For subsets  $A, B$  in  $P(X)$ ,  $A R B$  if and only if  $A \subset B$ . Is  $R$  an equivalence relation on  $P(X)$ ? Justify your answer.

**Solution:**

Since every set is a subset of itself,  $A \subset A$  for all  $A \in P(X)$ .

$\therefore R$  is reflexive.

Let  $ARB \Rightarrow A \subset B$

This cannot be implied to  $B \subset A$ .

If  $A = \{1, 2\}$  and  $B = \{1, 2, 3\}$ , then it cannot be implied that  $B$  is related to  $A$ .

$\therefore R$  is not symmetric.

If  $ARB$  and  $BRC$ , then  $A \subset B$  and  $B \subset C$ .

$\Rightarrow A \subset C$

$\Rightarrow ARC$

$\therefore R$  is transitive.

$R$  is not an equivalence relation as it is not symmetric.

**Question 9:**

Given a non-empty set  $X$ , consider the binary operation  $*$ :  $P(X) \times P(X) \rightarrow P(X)$  given by  $A * B = A \cap B \forall A, B$  in  $P(X)$  is the power set of  $X$ . Show that  $X$  is the identity element for this operation and  $X$  is the only invertible element in  $P(X)$  with respect to the operation  $*$ .

**Solution:**

$P(X) \times P(X) \rightarrow P(X)$  given by  $A * B = A \cap B \forall A, B$  in  $P(X)$

$A \cap X = A = X \cap A$  for all  $A \in P(X)$

$\Rightarrow A * X = A = X * A$  for all  $A \in P(X)$

$X$  is the identity element for the given binary operation  $*$ .

An element  $A \in P(X)$  is invertible if there exists  $B \in P(X)$  such that

$$A * B = X = B * A \quad [\text{As } X \text{ is the identity element}]$$

Or

$$A \cap B = X = B \cap A$$

This case is possible only when  $A = X = B$ .

$X$  is the only invertible element in  $P(X)$  with respect to the given operation  $*$ .

### Question 10:

Find the number of all onto functions from the set  $\{1, 2, 3, \dots, n\}$  to itself.

#### Solution:

Onto functions from the set  $\{1, 2, 3, \dots, n\}$  to itself is simply a permutation on  $n$  symbols  $1, 2, 3, \dots, n$ .

Thus, the total number of onto maps from  $\{1, 2, 3, \dots, n\}$  to itself is the same as the total number of permutations on  $n$  symbols  $1, 2, 3, \dots, n$ , which is  $n!$ .

### Question 11:

Let  $S = \{a, b, c\}$  and  $T = \{1, 2, 3\}$ . Find  $F^{-1}$  of the following functions  $F$  from  $S$  to  $T$ , if it exists.

i.  $F = \{(a, 3), (b, 2), (c, 1)\}$

ii.  $F = \{(a, 2), (b, 1), (c, 1)\}$

**Solution:**  $S = \{a, b, c\}, T = \{1, 2, 3\}$

i.  $F : S \rightarrow T$  is defined by  $F = \{(a, 3), (b, 2), (c, 1)\}$

$$\Rightarrow F(a) = 3, F(b) = 2, F(c) = 1$$

Therefore,  $F^{-1} : T \rightarrow S$  is given by  $F^{-1} = \{(3, a), (2, b), (1, c)\}$

ii.  $F : S \rightarrow T$  is defined by  $F = \{(a, 2), (b, 1), (c, 1)\}$

Since,  $F(b) = F(c) = 1$ ,  $F$  is not one-one.

Hence,  $F$  is not invertible i.e.,  $F^{-1}$  does not exist.

### Question 12:

Consider the binary operations  $*$ :  $R \times R \rightarrow R$  and  $\circ$ :  $R \times R \rightarrow R$  defined as  $a * b = |a - b|$  and  $a \circ b = a$ ,  $\forall a, b \in R$ . Show that  $*$  is commutative but not associative  $\circ$  is associative but not commutative. Further, show that  $\forall a, b, c \in R$ ,  $a * (b \circ c) = (a * b) \circ (a * c)$ . [ If it is so, we say that the operation  $*$  distributes over the operation  $\circ$ ]. Does  $\circ$  distribute over  $*$ ? Justify your answer.

#### Solution:

It is given that  $*$ :  $R \times R \rightarrow R$  and  $\circ$ :  $R \times R \rightarrow R$  defined as  $a * b = |a - b|$  and  $a \circ b = a$ ,  $\forall a, b \in R$ .

For  $a, b \in R$ , we have  $a * b = |a - b|$  and  $b * a = |b - a| = |-(a - b)| = |a - b|$

$$\therefore a * b = b * a$$

$\therefore$  The operation  $*$  is commutative.

$$(1 * 2) * 3 = (|1 - 2|) * 3 = 1 * 3 = |1 - 3| = 2$$

$$1 * (2 * 3) = 1 * (|2 - 3|) = 1 * 1 = |1 - 1| = 0$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3) \quad \text{where } 1, 2, 3 \in R$$

$\therefore$  The operation  $*$  is not associative.

Now, consider the operation  $\theta$ :

It can be observed that  $1\theta 2 = 1$  and  $2\theta 1 = 2$ .

$$\therefore 1\theta 2 \neq 2\theta 1 \quad (\text{where } 1, 2 \in R)$$

$\therefore$  The operation  $\theta$  is not commutative.

Let  $a, b, c \in R$ . Then, we have:

$$(aob)oc = aoc = a$$

$$ao(boc) = aob = a$$

$$\Rightarrow (aob)oc = ao(boc)$$

$\therefore$  The operation  $\theta$  is associative.

Now, let  $a, b, c \in R$ , then we have:

$$a * (boc) = a * b = |a - b|$$

$$(a * b)\theta(a * c) = (|a - b|)\theta(|a - c|) = |a - b|$$

$$\text{Hence, } a * (boc) = (a * b)\theta(a * c)$$

Now,

$$1\theta(2 * 3) = 1\theta(|2 - 3|) = 1\theta 1 = 1$$

$$(1\theta 2) * (1\theta 3) = 1 * 1 = |1 - 1| = 0$$

$$\therefore 1\theta(2 * 3) \neq (1\theta 2) * (1\theta 3) \quad \text{where } 1, 2, 3 \in R$$

$\therefore$  The operation  $\theta$  does not distribute over  $*$ .

### Question 13:

Given a non - empty set  $X$ , let  $*$ :  $P(X) \times P(X) \rightarrow P(X)$  be defined as  $A * B = (A - B) \cup (B - A)$ ,  $\forall A, B \in P(X)$ . Show that the empty set  $\Phi$  is the identity for the operation  $*$  and all the elements  $A$  of  $P(X)$  are invertible with  $A^{-1} = A$ .

(Hint:  $(A - \Phi) \cup (\Phi - A) = A$  and  $(A - A) \cup (A - A) = A * A = \Phi$ ).

### Solution:

It is given that  $*$ :  $P(X) \times P(X) \rightarrow P(X)$  is defined as  $A * B = (A - B) \cup (B - A)$ ,  $\forall A, B \in P(X)$   
 $A \in P(X)$  then,

$$A * \Phi = (A - \Phi) \cup (\Phi - A) = A \cup \Phi = A$$

$$\Phi * A = (\Phi - A) \cup (A - \Phi) = \Phi \cup A = A$$

$$\therefore A * \Phi = A = \Phi * A \quad \text{for all } A \in P(X)$$

$\Phi$  is the identity for the operation  $*$ .

Element  $A \in P(X)$  will be invertible if there exists  $B \in P(X)$  such that  
 $A * B = \Phi = B * A$  [As  $\Phi$  is the identity element]

$$A * A = (A - A) \cup (A - A) = \Phi \cup \Phi = \Phi \quad \text{for all } A \in P(X).$$

All the elements  $A$  of  $P(X)$  are invertible with  $A^{-1} = A$ .

### Question 14:

Define a binary operation  $*$  on the set  $\{0,1,2,3,4,5\}$  as

$$a + b = \begin{cases} a + b, & \text{if } a + b < 6 \\ a + b - 6 & \text{if } a + b \geq 6 \end{cases}$$

Show that zero is the identity for this operation and each element  $a \neq 0$  of the set is invertible with  $6 - a$  being the inverse of  $a$ .

### Solution:

Let  $X = \{0,1,2,3,4,5\}$

The operation  $*$  is defined as  $a + b = \begin{cases} a + b, & \text{if } a + b < 6 \\ a + b - 6, & \text{if } a + b \geq 6 \end{cases}$

An element  $e \in X$  is the identity element for the operation  $*$ , if  $a * e = a = e * a \quad \forall a \in X$

For  $a \in X$ ,

$$a * 0 = a + 0 = a \quad [a \in X \Rightarrow a + 0 < 6]$$

$$0 * a = 0 + a = a \quad [a \in X \Rightarrow 0 + a < 6]$$

$$\therefore a * 0 = a = 0 * a \quad \forall a \in X$$

Thus, 0 is the identity element for the given operation  $*$ .

An element  $a \in X$  is invertible if there exists  $b \in X$  such that  $a * b = 0 = b * a$ .

$$\text{i.e., } \left\{ \begin{array}{l} a + b = 0 = b + a, \quad \text{if } a + b < 6 \\ a + b - 6 = 0 = b + a - 6 \quad \text{if } a + b \geq 6 \end{array} \right\}$$

$$\Rightarrow a = -b \text{ or } b = 6 - a$$

$$X = \{0, 1, 2, 3, 4, 5\} \text{ and } a, b \in X. \text{ Then } a \neq -b.$$

$$\therefore b = 6 - a \text{ is the inverse of } a \text{ for all } a \in X.$$

Inverse of an element  $a \in X$ ,  $a \neq 0$  is  $6 - a$  i.e.,  $a^{-1} = 6 - a$ .

### Question 15:

Let  $A = \{-1, 0, 1, 2\}$ ,  $B = \{-4, -2, 0, 2\}$  and  $f, g: A \rightarrow B$  be functions defined by  $x^2 - x$ ,  $x \in A$  and

$$g(x) = 2 \left| x - \frac{1}{2} \right| - 1, \quad x \in A. \text{ Are } f \text{ and } g \text{ equal?}$$

### Solution:

It is given that  $A = \{-1, 0, 1, 2\}$ ,  $B = \{-4, -2, 0, 2\}$

Also,  $f, g: A \rightarrow B$  is defined by  $x^2 - x$ ,  $x \in A$  and  $g(x) = 2 \left| x - \frac{1}{2} \right| - 1, \quad x \in A.$

$$f(-1) = (-1)^2 - (-1) = 1 + 1 = 2$$

$$g(-1) = 2 \left| (-1) - \frac{1}{2} \right| - 1 = 2 \left( \frac{3}{2} \right) - 1 = 3 - 1 = 2$$

$$\Rightarrow f(-1) = g(-1)$$

$$f(0) = (0)^2 - 0 = 0$$

$$g(0) = 2 \left| 0 - \frac{1}{2} \right| - 1 = 2 \left( \frac{1}{2} \right) - 1 = 1 - 1 = 0$$

$$\Rightarrow f(0) = g(0)$$

$$f(1) = (1)^2 - 1 = 0$$

$$g(1) = 2 \left| 1 - \frac{1}{2} \right| - 1 = 2 \left( \frac{1}{2} \right) - 1 = 1 - 1 = 0$$

$$\Rightarrow f(1) = g(1)$$

$$f(2) = (2)^2 - 2 = 2$$

$$g(2) = 2 \left| 2 - \frac{1}{2} \right| - 1 = 2 \left( \frac{3}{2} \right) - 1 = 3 - 1 = 2$$

$$\Rightarrow f(2) = g(2)$$

$$\therefore f(a) = g(a) \quad \forall a \in A$$

Hence, the functions  $f$  and  $g$  are equal.

### Question 16:

Let  $A = \{1, 2, 3\}$ . Then number of relations containing  $(1, 2)$  and  $(1, 3)$  which are reflexive and symmetric but not transitive is,

- A. 1
- B. 2
- C. 3
- D. 4

### Solution:

The given set is  $A = \{1, 2, 3\}$ .

The smallest relation containing  $(1, 2)$  and  $(1, 3)$  which are reflexive and symmetric but not transitive is given by,

$$R = \{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 1), (3, 1)\}$$

This is because relation  $R$  is reflexive as  $\{(1, 1), (2, 2), (3, 3)\} \in R$ .

Relation  $R$  is symmetric as  $\{(1, 2), (2, 1)\} \in R$  and  $\{(1, 3), (3, 1)\} \in R$ .

Relation  $R$  is transitive as  $\{(3, 1), (1, 2)\} \in R$  but  $(3, 2) \notin R$ .

Now, if we add any two pairs  $(3, 2)$  and  $(2, 3)$  (or both) to relation  $R$ , then relation  $R$  will become transitive.

Hence, the total number of desired relations is one.

The correct answer is A.

### Question 17:

Let  $A = \{1, 2, 3\}$ . Then number of equivalence relations containing  $(1, 2)$  is,

- A. 1
- B. 2
- C. 3
- D. 4

### Solution:

The given set is  $A = \{1, 2, 3\}$ .

The smallest equivalence relation containing  $(1, 2)$  is given by;

$$R_1 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$$

Now, we are left with only four pairs i.e.,  $(2, 3), (3, 2), (1, 3)$  and  $(3, 1)$ .

If we add any one pair [say  $(2, 3)$ ] to  $R_1$ , then for symmetry we must add  $(3, 2)$ . Also, for transitivity we are required to add  $(1, 3)$  and  $(3, 1)$ .

Hence, the only equivalence relation (bigger than  $R_1$ ) is the universal relation.

This shows that the total number of equivalence relations containing  $(1, 2)$  is two.  
The correct answer is B.

### Question 18:

$$f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$$

Let  $f: R \rightarrow R$  be the Signum Function defined as  $f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$  and  $g: R \rightarrow R$  be the greatest integer function given by  $g(x) = [x]$ , where  $[x]$  is greatest integer less than or equal to  $x$ . Then does  $f \circ g$  and  $g \circ f$  coincide in  $(0, 1]$ ?

### Solution:

$$f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$$

It is given that  $f: R \rightarrow R$  be the Signum Function defined as

Also  $g: R \rightarrow R$  is defined as  $g(x) = [x]$ , where  $[x]$  is greatest integer less than or equal to  $x$ .

Now let  $x \in (0, 1]$ ,

$$[x] = 1 \text{ if } x = 1 \text{ and } [x] = 0 \text{ if } 0 < x < 1.$$

$$\therefore fog(x) = f(g(x)) = f([x]) = \begin{cases} f(1), & \text{if } x = 1 \\ f(0), & \text{if } x \in (0,1) \end{cases} = \begin{cases} 1, & \text{if } x = 1 \\ 0, & \text{if } x \in (0,1) \end{cases}$$

$$\begin{aligned} g \circ f(x) &= g(f(x)) \\ &= g(1) \quad [x > 0] \\ &= [1] = 1 \end{aligned}$$

Thus, when  $x \in (0,1)$ , we have  $fog(x) = 0$  and  $gof(x) = 1$ .

Hence,  $fog$  and  $gof$  does not coincide in  $(0,1]$ .

### Question 19:

Number of binary operations on the set  $\{a,b\}$  are

- A. 10
- B. 16
- C. 20
- D. 8

### Solution:

A binary operation  $*$  on  $\{a,b\}$  is a function from  $\{a,b\} \times \{a,b\} \rightarrow \{a,b\}$

i.e.,  $*$  is a function from  $\{(a,a), (a,b), (b,a), (b,b)\} \rightarrow \{a,b\}$

Hence, the total number of binary operations on the set  $\{a,b\}$  is  $2^4 = 16$ .

The correct answer is B.